

I GRAPHS

I.1. The Basics

1.1. Recall

1) A set is merely an accumulation of objects.

These objects are called **elements** of the set. If an object x is an element of S , we write $x \in S$. The set of all elements with a certain property P is denoted via $\{x \mid x \text{ has property } P\}$.

2) An **n-ary relation** R on a set A is a subset of the power set of A^n , i.e. $R \subseteq P(A^n)$. If $n=2$, we call the relation **binary**.

A binary relation R on a set A is called

i) **symmetric** if $R(a,b) \Rightarrow R(b,a)$ for all $a, b \in A$

ii) **asymmetric** if $R(a,b) \Rightarrow \neg R(b,a)$ for all $a, b \in A$.

iii) **antisymmetric** if $R(a,b), R(b,a) \Rightarrow a = b$ for all $a, b \in A$.

iv) **reflexive** if $R(a,a)$ for all $a \in A$

v) **irreflexive** if $\neg R(a,a)$ for all $a \in A$

vi) **transitive** if $R(a,b) \wedge R(b,c) \Rightarrow R(a,c)$ for all $a, b, c \in A$.

1.2 Definition

A graph $G = (V, E)$ is a pair of sets V and E st. E consists of subsets of V of size two. V is called the set of vertices and E the set of edges. A graph G is called finite if V is a finite set. The order $|G|$ of a graph $G = (V, E)$ is the cardinality of its vertex set, so $|G| = |V|$. The size $\|G\|$ of G is the cardinality of its edge set, $\|G\| = |E|$.


1.3 Visualisation

Let $G = (V, E)$ be a graph. We visualise vertices $u, v, \dots \in V$ by dots and edges $e = \{u, v\} \subset E$ by the diagram



1.4 Example

Let $G = (V, E)$ be the graph with $V = \{a, b, c, d, e\}$ and $E = \{\{a, b\}, \{a, c\}, \{a, d\}, \{a, e\}, \{b, c\}, \{d, e\}\}$. The G has order 5 and size 6. It can be visualized via



This visualisation motivates its name: **bowtie graph**.

1.5 Notation

- 1) For a graph $G = (V, E)$ we may denote its vertex set by $V(G)$ or V_G for clarity.
- 2) Similarly, we often denote E by $E(G)$ or E_G .
- 3) We denote an edge $\{u, v\}$ simply by uv .
- 4) Edges are often called e, e_1, e_2, f, \dots , while vertices are called u, v, x, y, \dots

1.6 Definition

Let $G = (V, E)$ be a graph.

- 1) If $uv \in E$ is an edge, then we say that u and v are adjacent or neighbours. If $uv \notin E$, we call u and v nonadjacent.
- 2) If $e = uv \in E$, we say that u and v are the end vertices of e or that they are incident with e .
- 3) The neighborhood $N(v)$ of a vertex $v \in V$ is the set of all vertices adjacent to v , i.e. $N(v) := \{u \in V \mid uv \in E\}$. The closed neighborhood $N[v]$ of v is $N[v] := N(v) \cup \{v\}$. (Think of (a, b) vs. $[a, b]$)
- 4) The neighborhood $N(S)$ of a set of vertices is defined as $N(S) := \bigcup_{v \in S} N(v)$. Similarly, the closed neighborhood $N[S]$ is set to be $N[S] := N(S) \cup S$ ($= \bigcup_{v \in S} N[v]$).
- 5) The degree $\deg(v)$ of $v \in V$ is the number of edges incident with v , i.e. $\deg(v) := |\{e \in E \mid v \in e\}| = |N(v)|$.

6) The maximum degree $\Delta(G)$ of G is defined as

$$\Delta(G) := \max \{ \deg(v) \mid v \in V \}. \text{ Similarly, } \delta(G) := \min \{ \deg(v) \mid v \in V \}$$

is the minimum degree of G .

7) The degree sequence of a graph G is the sequence containing all degrees of the vertices of G (with repetition) in decreasing order.

1.7 Example


Consider G given by

$$\text{Then } \Delta(G) = 5$$

$$\delta(G) = 1$$

$$N(c) = \{a, d, f, g\}$$


$$N(b) = \{b, d\} \quad N[a, g] = \{a, c, d, e, f, g\}$$



order of G is 7, size of G is 9. Degree sequence is $(5, 3, 3, 2, 2, 2, 1)$.

1.8 Example

Consider G given via the diagram



$$\text{Then } V(G) = \{a, b, c, d, e, f, g, h, i\}$$

$$\bar{E}(G) = \{\{a, d\}, \{a, e\}, \{b, c\}, \{b, d\}, \{b, g\}, \{c, f\}, \{d, f\}, \{d, g\}, \{g, h\}\}$$

order $|G| = 9$, size of G is 9, degree sequence $(3, 3, 3, 2, 2, 2, 1, 0)$,

$$N(f) = \{c, d\}, \quad N[d, e] = \{a, b, c, d, e, f\}, \quad \Delta(G) = 3, \quad \delta(G) = 0$$

What is the sum?


1.9 Remark

A graph can be considered as a set V together with a binary relation E on V which is symmetric and irreflexive.

1.10 Definition (Variants of Graphs)

1) If $G = (V, E)$ and we replace E with a set of ordered pairs, then we call G a **directed graph**, or **digraph**.

Ex:




$$V(G) = \{a, b, c, d\}$$

$$E(G) = \{(a, b), (a, c), (a, d), (d, a), (d, c)\}$$

2) If $G = (V, E)$ and we replace E by a multiset, iterations of the same elements are distinguished, then we call G a **multigraph**.


Ex



$$E = [\{a, b\}, \{b, c\}, \{b, c\}, \{b, c\}]$$

3) If $G = (V, E)$ and we extend E by allowing loops, we call G a **pseudograph**.


Ex



$$E = \{\{a, b\}, \{b, d\}, [c, c], [d, d]\}.$$

4) If we allow our edges to be arbitrary sets of vertices instead of 2-elementary ones, we call G a **hypergraph**.

Ex:



$$E = \{\{a\}, \{b\}, \{a, b, d\}, \{c, d\}\}.$$

1.11 Setting

In this lecture, unless otherwise stated, by a graph we mean a finite, simple graph with $|V| \geq 1$.

1.12 Definition

- The **Complete graph** K_n for $n \geq 1$ is the graph consisting of n vertices such that any two vertices are adjacent.

e.g. K_1 K_2 K_3 K_4 K_5 ...
• •—•   

- The **empty graph** E_n is the graph consisting of n vertices and no edges, e.g. E_1 E_2 E_3 E_4 ...
• •—• •—• •—• •—•

INTERMEZZO - 20 min

- 1) Illustrate your given scenario with a graph. Determine size, order, degree sequence, $\Delta(G)$ and $\delta(G)$. Compare $|E|$ and sum of degrees.
- 2) Formulate as many theorems about complete graphs as you can.

- (A) V consists of all students of groups A, B and C. Students x and y are connected, if they share a common (Arabic) letter in their first name.
- (B) V consists of all students of groups B, C and D. Students x and y are connected, if they have attended the same course (same time + section) before this term.
- (C) V consists of all students of groups C, D and E.. Students x and y are connected, if they share a common declared major.
- (D) V consists of all students of groups D, E and A . Students x and y are connected, if they come from the same city and district.
- (E) V consists of all students of groups E, A and B . Students x and y are connected, if they attend no more than one common course this term.

1.13 The Handshaking Lemma

If $G = (V, E)$ is a graph, then

$$\sum_{v \in V} \deg(v) = 2|E|. \quad (*)$$

Proof

We proceed by induction on $n := |E|$.

$n=0$: If $|E|=0$, then $\deg(v)=0$ for any $v \in V$; whence clearly.

$$0 = \sum_{v \in V} \deg(v) = 2|E|=0.$$

$n \mapsto n+1$: Assume $(*)$ holds for any $G' = (V', E')$ with $|E'| = n$ (I.H.) and

consider $G = (V, E)$ with $|E| = n+1 (\geq 1)$ arbitrary. Let $e \in E$

arbitrary and consider $G' = (V, E \setminus \{e\})$. Then, if $e = uv$,

we get $|E(G)| = |E(G')| + 1$ and

$$\deg(u) = \deg^{G'}(u) + 1 \text{ and } \deg(v) = \deg^{G'}(v) + 1, \text{ whence}$$

$$2|E(G)| = 2|E(G')| + 2$$

$$\begin{aligned}
 &\stackrel{\text{I.H.}}{=} \sum_{w \in V} \underbrace{\deg^{G'}(w)}_{\deg^{G'}(x)} + 2 \\
 &= \sum_{w \in V \setminus \{u, v\}} \underbrace{\deg^{G'}(w)}_{\deg^{G'}(u)} + \underbrace{\deg^{G'}(u) + 1}_{\deg^{G'}(v)} + \underbrace{\deg^{G'}(v) + 1}_{\deg^{G'}(v)} \\
 &= \sum_{w \in V} \underbrace{\deg^{G'}(w)}_{\deg^{G'}(w)}, \text{ as desired.} \quad \blacksquare
 \end{aligned}$$

1.14 Corollary

Any graph G has an even number of vertices of odd degree.

Proof - Exercise.

1.15 Corollary

For any graph $G = (V, E)$ we have

$$\delta(G) \leq 2 \frac{|E|}{|V|} \leq \Delta(G).$$

Proof

$$|V| \cdot \delta(G) = \sum_{v \in V} \delta(G) \leq \sum_{v \in V} \deg(v) \leq \sum_{v \in V} \Delta(G) = |V| \Delta(G)$$

|| 1.13

$$2|E|.$$

$$\Rightarrow \delta(G) \leq 2 \frac{|E|}{|V|} \leq \Delta(G), \text{ as desired.}$$

1.16 Lemma

If $|G| \geq 2$, then G contains at least two vertices of the same degree.

Proof

If G has two vertices of degree 0, then we are done.

Otherwise, we may assume that G has none. If $|G|=n$, and $v \in V$, then $1 \leq \deg(v) \leq n-1$. Note that this leaves us with $n-1$ choices of degrees for n many different vertices. Hence, at least two vertices must have the same degree. \blacksquare


1.17 Remark

The above line of thought is called the **pigeon hole principle**.

If there are n many pigeons wanting to fit into $n-1$ many holes, then at least two of them have to cuddle up in the same hole.


1.18 Definition

- i) The path P_n is the graph on n vertices v_1, \dots, v_n with the edge set $E(P_n) = \{v_i v_{i+1} \mid 1 \leq i < n\}$, i.e. P_n is represented by the diagram




- ii) The cycle C_n the graph on n vertices with the edge set

$$E(C_n) = \{v_i v_{i+1} \mid 1 \leq i < n\} \cup \{v_n v_1\}. \text{ Eg.}$$



- iii) Let $G = (V, E)$ be an arbitrary graph. The complement \bar{G} of G is the graph $\bar{G} = (\bar{V}, \bar{E})$, where $\bar{E} = \{uv \mid u, v \in V, uv \notin E\}$.

e.g.



1.19 Definition


We call a graph G regular if any of its vertices has the same degree. If this degree is r , we say that G is r -regular.

1.20 Remarks

- i) A graph G is regular iff $\delta(G) = \Delta(G)$.
- ii) P_n is $(n-1)$ -regular and E_n is 0-regular.
- iii) A r -regular graph of order n has $\frac{1}{2}nr$ many edges.

1.21 Example

The graph below is 4-regular of order 6.



1.4 Subgraphs

There are two ways in which one graph can be part of another graph.

1.22 Definition

- 1) A graph H is called a **subgraph** of some graph G , write $H \subseteq G$, if $V(H) \subseteq V(G)$ and $E(H) \subseteq E(G)$. (possibly after labelling the vertices in G). We also say that G **contains** H .
- 2) If $H \subseteq G$, we say that H is an **induced subgraph** of G , write $H \preceq G$, if $E(H) = \{uv \in E(G) \mid u, v \in V(H)\}$.







1.23 Remark

- i) $H \subseteq G$ is induced if for any two vertices in H we have:
If they are adjacent in G , then they are adjacent in H .
- ii) Every induced subgraph is a subgraph but not vice versa.
- iii) If G is a graph and $S \subseteq V(G)$, then there is only one induced subgraph $H \preceq G$ with vertex set S , i.e. $V(H)=S$.
we denote this graph by $\langle S \rangle$ and call it the subgraph of G induced by S .

1.24 Example

Consider G given as

Then



subgraph ✓ ✓ ✗ ✓ ✓

induced ✗ ✓ ✗ ✓ ✗

7.5 Walks in Graphs

1.25 Definition

A (v_0, v_k) -walk in a graph is a sequence of vertices (v_0, v_1, \dots, v_k) s.t. any two consecutive vertices v_i and v_{i+1} are adjacent. We call the edges $\{v_0v_1, v_1v_2, \dots, v_{k-1}v_k\}$ the edges of the walk. We say that the walk is **Closed** if $v_0 = v_k$. The **length** of a walk is the number of edges in it, counting repetition. Here, the length is k .


1.26 Definition

We distinguish the following types of walks:

- A **trail** is a walk whose edges are pairwise distinct.
- A **circuit** is a closed walk whose edges are pw. distinct.
- A **path** is a walk whose vertices are distinct.
- A **cycle** is a closed walk $(v_0, v_1, \dots, v_k, v_0)$ with $k \geq 2$ and whose vertices v_0, \dots, v_k are pairwise distinct.

1.27 Example

Consider G via



Give examples for a

- walk (d, b, c, d, b, a)

da-walk, length 5

- trail (d, c, a, b, c, e)

de-trail, length 5

- path (d, c, a, b, e)

de-path, length 4

- closed walk (e, b, c, a, b, d, e)

e-closed walk, length 6

- circuit (d, c, a, b, c, e, d)

d-circuit, length 6

- cycle (d, c, a, b, c, d)

d-cycle, length 5


INTERMEZZO ~ 15 min

1) Draw the Venn-Diagram describing the relation between:

trails, finite sequences of vertices, paths,
closed walks, cycles, walks and circuits.

2) Consider the graph G below. Place the sequences below into your diagram.

$\bullet G =$



$s_1 = (g, d, b, c, a, b, g)$ circuit

$s_2 = (a, c, f, c, a, d)$ walk

$s_3 = (a, b, c, a)$ cycle

$s_4 = (b, a, c, b, d)$ trail

$s_5 = (f, c, a, b, e)$ nothing

$s_6 = (d, g, b, a, c, f, e)$ path

$s_7 = (g, d, f, e, d, g, b, g)$ closed walk

1.28 Lemma

If $\delta(G) \geq 2$, then G contains a cycle as a subgraph.

Proof

Let $p = (v_0, v_1, \dots, v_k)$ be a path in G of maximal length.

This exists, as G is finite. Further, as $\delta(G) \geq 2$, we get $k \geq 2$.

As $\deg(v_0) \geq \delta(G) \geq 2$, v_0 has at least two neighbors. One of them is v_1 . Let us denote the other one by u . If $u \neq v_i$ for all $1 \leq i \leq k$, then $\tilde{p} = (u, v_0, v_1, \dots, v_k)$ is still a path and of greater length than p , contradicting our assumptions.

Hence, $u = v_i$ for some $1 \leq i \leq k$. But then the sequence

$(v_0, v_1, \dots, v_{i-1}, v_i = u, v_0)$ is the desired cycle subgraph of G .

1.29 Corollary (Contrapositive)

If G does not contain any cycles, then $\delta(G) \leq 1$.

1.30 Theorem

Every uv -walk in a graph contains a uv -path.

Proof

We proceed by strong induction on the length n of the walk.

1.B $n=1$. If the uv -walk is of length one, then it is exactly

(u, v) , which is also a path.

1.5. Assume every uv -walk of length at most $n \geq 1$ contains a uv -path. (I.H.)

Assume there is a uv -walk $w = (u = w_0, w_1, w_2, \dots, w_n, w_{n+1} = v)$ of length $n+1$. If w is already a path, we are done. Otherwise there are i, j st. $0 \leq i < j \leq n+1$ and $w_i = w_j$. But then the walk \tilde{w} which arises from w by deleting the vertices $w_{i+1}, \dots, w_{j-1}, w_j$, i.e. $\tilde{w} = (u = w_0, w_1, \dots, w_i, w_{j+1}, \dots, w_n = v)$ is still a uv -walk, but of length at most n . Using I.H., we know that \tilde{w} contains a uv -path, whence also w contains (the same) uv -path. □

1.6 Connectivity

1.31 Definition

A graph is **connected** if there exists an uv -path in G for any vertices $u, v \in V(G)$. Otherwise, it is called **disconnected**.

Intuition

A graph is connected if you could pick it up entirely by just lifting one vertex. If it is not connected, then the subgraph you lift that way is called a **connected component**.

1.32 Definition

A **connected component** of G is a maximal connected induced subgraph of G . I.e. $C \subseteq G$ is a connected component iff i) C is connected and ii) for any $v \in V(G) \setminus V(C)$ the induced subgraph on $V(C) \cup \{v\}$ is not connected.

1.33 Remark

G is connected iff it has exactly one connected component.

Even among connected graphs, there different levels of being connected. E.g. the graph K_5  "feels" more connected than the graph . In order to properly describe this intuition, we need more notations.


1.34 Definition (Vertex and Edge Deletion)


Let G be a graph, $S \subseteq V_G$, and $T \subseteq E_G$.

- 1) By $G-S$ we denote the graph arising from G by removing from V_G all vertices in S and their incident edges..
- 2) If $S = \{v\}$ is one vertex, we also write $G-v$ for $G-\{v\}$.
- 3) By $G-T$ we denote the graph arising from G by removing only the edges in T , but no vertices.
- 4) If $T = \{e\}$ is only one edge, we also write $G-e$ for $G-\{e\}$.


1.35 Example

Consider G as given below. Note that G only has one connected component.




Then $G-d$ is  and has 2 connected components.

The vertex d is called a **cut vertex**.

Further, $G-\{e,f\}$ is . It also has 2 connected components.

The set $\{e,f\}$ is called a **cut set**.

Further, $G-ab$ is  again, it has 2 connected components.

We call the edge ab a **bridge**.

1.36 Definition

Let G be a graph.

- i) We call $v \in V_G$ a **cut vertex** if $G-v$ has more connected components than G itself.
- ii) We call $e \in E_G$ a **bridge** if $G-e$ has more connected components than G itself.
- iii) We call $S \subseteq V_G$ a **cut set** if $G-S$ is disconnected.
- iv) A connected graph which does not contain any cut vertices is called **non-separable**.

1.37 Observation

- 1) If G is connected then v is a cut vertex of G iff $\{v\}$ is a cut set.
- 2) The vertex v is a cut vertex iff there are vertices u and w , different from v s.t. every uw -path uses v .
- 3) A graph has no cut sets iff it is a complete graph. (Hw)


1.38 Definition

For a non-complete graph G , we define its **connectivity** $\kappa(G)$

as the minimal size of a cut set. For K_n , we set $\kappa(K_n) := n-1$.

INTERMEZZO

- 1) Consider the graph G below. Find all cut vertices, bridges, cut sets which do not contain proper cut subsets and $\kappa(G)$.



- 2) Is there any impact on $\deg(v)$ on whether or not v can be a cut vertex?
- 3) If S is a cut set which does not contain a proper cut subset, what do you observe for S ? Can you prove your observation?

1.39 Lemma

If G is a nonseparable graph of order at least 3, then $\delta(G) \geq 2$ and every vertex of G is contained in a cycle.

Proof


Consider G nonseparable with $|G| \geq 3$. By definition, G is connected, i.e. $\delta(G) \geq 1$.

First we show that $\delta(G) \geq 2$. Otherwise, we have $\delta(G) = 1$, i.e. there is some vertex v s.t. $\deg(v) = 1$. Let u be the unique neighbor of v and w any other vertex of G (which exists as $|G| \geq 3$). Then clearly any vw -path must use the unique neighbor u of v ; whence u is a cut vertex. This contradicts the fact that G is inseparable. Hence, $\delta(G) \geq 2$, as desired.

Picture:



Now, consider $v \in V_G$ arbitrary. We want to show that v is contained in a cycle in G . As $\delta(G) \geq 2$, v has at least 2 neighbors, say u and w . As G is nonseparable, $G - v$ is still connected. In particular, there is a uw -path ($u = x_0, x_1, x_2, \dots, x_\ell = w$) in $G - v$. But then the walk $(x_0 = u, x_1, \dots, x_\ell = w, v, x_0 = u)$ is the desired cycle containing v .



1.40 Definition

We say that G is k -connected if $\kappa(G) \geq k$, i.e. if G is connected and $G - S$ is still connected for any $S \subseteq V_G$ with $|S| < k$.

1.41 Lemma

The following hold.

- 1) G is connected iff $\kappa(G) \geq 1$.
- 2) G is 1-connected iff G is connected.
- 3) G is 2-connected iff G is connected and has no cut vertices.
- 4) G is 2-connected iff G is non-separable.
- 5) If G is 2-connected, then it contains at least one cycle.
(for $|G| \geq 3$)
- 6) If G is k -connected, then G is j -connected for all $j \leq k$.
- 7) $|G| > \kappa(G)$.
- 8) $\kappa(G) \leq \delta(G)$.

Proof

1)-6) are easy observations - verify them by yourselves.

7) If $G = K_n$, then $|G| = n > n-1 = k(G)$. Otherwise, assume $k(G) = \ell$, i.e.

ex. $S \subseteq V_G$ s.t. $|S| = \ell$ and $G-S$ is disconnected. For $G-S$ to be disconnected, it must contain at least 2 vertices, whence

$$|G| \geq |S| + 2 = k(G) + 2 > k(G).$$

8) Assume $k(G) > \delta(G)$ and let $v \in V_G$ s.t. $\deg(v) = \delta(G)$. Note that $|G| > k(G) > \delta(G) = |N(v)|$, whence $G - N(v)$ contains at least one vertex besides G . But clearly, $G - N(v)$ is disconnected, as $\deg(v) = 0$.

Hence, $N(v)$ is a cut set and $k(G) \leq |N(v)| = \delta(G)$, contradicting the assumptions. □

1.7 Bipartite Graphs


1.42 Definition

A graph G is called **bipartite** if we can partition the vertex set V_G into two disjoint sets $V_G = X \cup Y$ s.t. every edge of G has one end vertex in X and the other in Y .

1.43 Example

Consider

$$G :=$$




We can partition the vertices of G

into two sets via


$$X = \{x_1, x_2, x_3\},$$

$$Y = \{y_1, y_2, y_3, y_4\}.$$



Rearranging the position of the vertices makes it clear that

G is bipartite:



We see that there are no edges between any two vertices in X or in Y .

1.44 Remark


Following the above we see that a graph G is bipartite if and only if we can color the vertices of G with two colors s.t. the end vertices of each edge have different colors.

1.45 Definition


Let $m, n \in \mathbb{Z}_+$. The complete bipartite graph $K_{m,n}$ is the bipartite graph with $X = \{x_1, \dots, x_m\}$, $Y = \{y_1, \dots, y_n\}$, $V_G = X \cup Y$ and $E_G = \{xy \mid x \in X, y \in Y\}$.

1.46 Example


Below are some examples of complete bipartite graphs.



$K_{1,3}$



$K_{4,2}$



$K_{3,3}$

The following theorem helps us decide whether or not a given graph is bipartite.

1.47 Theorem

A graph is bipartite iff it does not contain odd cycles.

Proof

" \Rightarrow " Assume G is bipartite and nevertheless there is a cycle of odd length, say $(x_0, x_1, x_2, \dots, x_{2k}, x_{2k+1} = x_0)$. By

Remark 1.44, we can color V_G in two colors, C_1 and C_2 , s.t. adjacent vertices have different colors. Then, if

x_0 has color C_1 , x_1 has color C_2 whence x_2 has color C_1 .

That way we see that the color of x_i is $\begin{cases} C_1 & \text{if } i \text{ is even} \\ C_2 & \text{if } i \text{ is odd.} \end{cases}$

Following that logic, the vertex $x_0 = x_{2k+1}$ should have

color C_1 and color C_2 at the same time, which is a contradiction.

" \Leftarrow " Now consider that G does not contain odd cycles.

We will show that G is bipartite by providing a partition.

We may assume that G is connected as otherwise we work component per component.

Pick $v \in V_G$ arbitrary and define

$X = \{w \in V_G \mid \text{the shortest } vw \text{ path has even length}\}$ and

$Y = \{w \in V_G \mid \text{the shortest } vw \text{ path has odd length}\}$.

Clearly, X and Y are disjoint. We will show that there are no adjacent vertices in X or Y respectively.

Note that $v \in X$.

aiming for a contradiction, assume that there are vertices $w_1, w_2 \in X$ which are adjacent. Clearly, $w_i \neq v$, as otherwise the shortest vw_2 -path was exactly vw_2 of length 1. Similarly, $w_2 \neq v$. Let $p_1 = (v = x_0, x_1, \dots, x_{2k} = w_1)$ and $p_2 = (v = y_0, y_1, \dots, y_{2l} = w_2)$ be the shortest vw_1 -and vw_2 -pathes.

Suppose that $x_i = y_j$ for some $0 < i \leq 2k$ and $0 < j \leq 2l$.

If $i < j$, then $(v = x_0, x_1, \dots, x_i, y_{j+1}, \dots, y_{2l} = w_2)$ is a vw_2 path shorter than p_2 , a contradiction. Similarly, $j < i$ is impossible, whence $i = j$, whenever $x_i = y_j$.

Now, pick the largest i s.t. $x_i = y_i$. As $x_0 = v = y_0$, such an i always exists. Then we obtain the following

cycle $C = (\underbrace{x_i, x_{i+1}, \dots, x_{2k}}_{2k-i} = w_1, \underbrace{w_2 = y_{2l}, y_{2l-1}, \dots, y_{i+1}}_1, \underbrace{y_i = x_i}_{2l-i})$.

This is a cycle, as p_1 and p_2 were paths and i was maximal s.t. $x_i = y_i$. Further, the length of C is odd, as it equals $(2k-i) + 1 + (2l-i) = 2(k+l-i) + 1$.


This contradicts our assumption that G does not contain odd cycles. We hence proved that no two vertices w_1 and w_2 from X can be adjacent.

The arguments for $v_1, v_2 \in Y$ is analogous (try to write it down).

This concludes the proof. 

1.8 Graph Isomorphisms


In Example 1.43, we rearranged the given graph




We understand G_1 and G_2 as the same, even though on the first glance, they look very similar. Another example is

given by

$$G =$$



and



We can relabel the vertices of G via $a \mapsto 1, b \mapsto 2, c \mapsto 8, d \mapsto 3, e \mapsto 7, f \mapsto 4, g \mapsto 6$ and $h \mapsto 5$ and obtain H . The aim of this section is to formalise this concept.

1.48 Definition

We say that a graph G is **isomorphic** to a graph H if there exists a bijection $\varphi: V_G \rightarrow V_H$ s.t. for any $u, v \in V_G$ we have that $\{u, v\} \in E_G$ if and only if $\{\varphi(u), \varphi(v)\} \in E_H$. Then, the map φ is called an **isomorphism**. And we write $G \cong H$.

1.49 Remark

Let $G \cong H$ via $\varphi: V_G \rightarrow V_H$. Then the following hold.

1) $|V_G| = |V_H|$ and $|E_G| = |E_H|$ and $\bar{G} \cong \bar{H}$.

2) The degree sequence of G equals the degree sequence of H .



3) G is connected iff H is connected.

4) $\deg^G(v) = \deg^H(\varphi(v))$ for all $v \in V_G$.







INTERMEZZO

1) Compute $|G_i|$, $\|G_i\|$ and degree sequence of G_i for $i=1,2$.

Is $G_1 \cong G_2$? Why?



2) Are the 2 graphs below isomorphic? Argue!



Chapter 2 - Distance in Graphs

We have a natural understanding of the "distance" between two objects in our physical space. But there are many other ways of defining distances. E.g., the distance between two people could be the positive difference of their birth years or the number of acquaintances you need to connect one to the other.

In this chapter we will introduce a notion of distance of vertices in a graph. But first let us note what are the characterising properties that make us call all these concepts "distances".

2.1 Definition


Let X be any set. We call a function $d: X \times X \rightarrow \mathbb{R}^{\geq 0}$ a **metric** if it satisfies for all $x, y, z \in X$:

1) $d(x, y) \geq 0$

2) $d(x, y) = 0$ iff $x = y$

3) $d(x, y) = d(y, x)$

4) $d(x, z) \leq d(x, y) + d(y, z)$ Triangle Inequality



We then call the pair (X, d) a **metric space**.

2.2 Example

Consider $X = \mathbb{R}$ and $d: \mathbb{R} \times \mathbb{R} \rightarrow \mathbb{R}^{>0}$ via

$d(x,y) := |x-y|$. Then (\mathbb{R}, d) is a metric space.

Now we are ready to define a metric on an arbitrary graph.

2.3 Definition

Let G be any graph and $u, v \in V_G$. We define the distance $d(uv)$ between u and v as the length of the shortest uv -path in G , i.e.

$$d(uv) = \min \{ \text{length}(p) \mid p \text{ is a } uv\text{-path} \}.$$

If there is no such path, we set $d(uv) := \infty$.

2) If $d(uv) = k$, then any uv -path of length k is called a **geodesic**.

2.4 Remark

1) We may write $d^G(uv)$ to emphasize that we consider the distance in G .


2) While in (\mathbb{R}, d) geodesics are unique, in general this is not the case. Consider for example two opposite poles on a sphere.

3) $d(xy) = \infty$ iff x and y are in different connected components.

4) (G, d) is a metric space for any connected graph G .

We call something eccentric if it is away from the usual.

Similarly, in graphs we measure by eccentricity how far a vertex is from the center. Consider the following notions on a cycle:



2.5 Definition

- 1) The eccentricity $\text{ecc}(v)$ of a vertex v is its greatest distance to any other vertex, i.e. $\text{ecc}(v) = \max \{ d(v, u) \mid u \in V_G \}$.
- 2) The radius $\text{rad}(G)$ is the smallest possible eccentricity and the diameter $\text{diam}(G)$ is the largest possible eccentricity.
- 3) The center $C(G)$ is the set $\{ v \in V_G \mid \text{ecc}(v) = \text{rad}(G) \}$ and the periphery $P(G)$ is the set $\{ v \in V_G \mid \text{ecc}(v) = \text{diam}(G) \}$.

2.6 Example

- 1) Consider P_5 , the path of length 4, i.e.

Then $d(v_i, v_j) = |i-j|$ whence $\text{ecc}(v_i) = \max \{ 0, 1, 2, 3, 4 \} = 4$.
 $d(v_2, v_i) = |i-2|$, whence $\text{ecc}(v_2) = \max \{ 1, 0, 1, 2, 3 \} = 3$
 $d(v_3, v_i) = |i-3|$, whence $\text{ecc}(v_3) = \max \{ 2, 1, 0, 1, 2 \} = 2$
 $d(v_4, v_i) = |i-4|$, whence $\text{ecc}(v_4) = \max \{ 3, 2, 1, 0, 1 \} = 3$
 $d(v_5, v_i) = |i-5|$, whence $\text{ecc}(v_5) = \max \{ 4, 3, 2, 1, 0 \} = 4$.


Hence $\text{rad}(P_5) = \min \{ \text{ecc}(v) \mid v \in V \} = \min \{ 4, 3, 2, 3, 4 \} = 2$.

Also $C(P_5) = \{ v \in V \mid \text{ecc}(v) = \text{rad}(P_5) \} = \{ v_2 \}$.

Further $\text{diam}(P_5) = \max \{ \text{ecc}(v) \mid v \in V \} = \max \{ 4, 3, 2, 3, 4 \} = 4$.

and $P(P_5) = \{ v \in V \mid \text{ecc}(v) = \text{diam}(P_5) \} = \{ v_0, v_4 \}$.

2) Consider $G := C_6$, the cycle of length 6,



$$\text{Then } d(v_0, v_i) = |3 - i|, \text{ecc}(v_0) = \max\{0, 1, 2, 3, 2, 1\} = 3,$$

$$d(v_1, v_i) = |3 - 1 - i|, \text{ecc}(v_1) = \max\{1, 0, 1, 2, 3, 2\} = 3$$

$$d(v_2, v_i) = |3 - 1 - 1 - i|, \text{ecc}(v_2) = \max\{2, 1, 0, 1, 2, 3\} = 3$$

$$d(v_3, v_i) = |3 - i|, \text{ecc}(v_3) = \max\{3, 2, 1, 0, 1, 2\} = 3$$

$$d(v_4, v_i) = |3 - 1 - i|, \text{ecc}(v_4) = \max\{2, 3, 2, 1, 0, 1\} = 3$$

$$d(v_5, v_i) = |3 - 1 - 1 - i|, \text{ecc}(v_5) = \max\{1, 2, 3, 2, 1, 0\} = 3.$$

$$\text{Hence, } \text{rad}(G) = \min\{\text{ecc}(v) \mid v \in V_G\} = \min\{3, 3, 3, 3, 3, 3\} = 3$$

$$\text{whence } C(G) = \{v \in V_G \mid \text{ecc}(v) = \text{rad}(G)\} = V_G.$$

$$\text{Further, } \text{diam}(G) = \max\{\text{ecc}(v) \mid v \in V_G\} = \max\{3, 3, 3, 3, 3, 3\} = 3.$$

$$\text{and } P(G) = \{v \in V_G \mid \text{ecc}(v) = \text{diam}(G)\} = V_G.$$


INTERMEDIATE

I: Find $\text{rad}(G)$, $\text{diam}(G)$, $C(G)$ and $P(G)$ of the following:

$$1) G_1 = \emptyset_{10}$$

$$2) G_2 = K_6$$

II Consider



Find:

$$- d(b, c), d(b, k), d(c, m)$$

$$- \text{ecc}(v) \text{ for all } v \in V_G$$

$$- \text{rad}(G), \text{diam}(G), C(G), P(G).$$

2.7 Lemma

For any graph G we have $\text{rad}(G) \leq \text{diam}(G) \leq 2 \text{rad}(G)$.

Proof

We have $\text{rad}(G) \leq \text{diam}(G)$ by definition. For the other inequality, pick $v \in C(G)$ arbitrary and consider $u, w \in V_G$ arbitrary s.t. $d(u, w) = \text{diam}(G)$. Then


$$d(u, w) \leq d(u, v) + d(v, w) \leq \text{ecc}(v) + \text{ecc}(v) = 2 \text{rad}(G). \quad \blacksquare$$

2.8 Theorem

Every graph G is isomorphic to the graph induced by the center of another graph H , i.e. ex. H s.t. $G \cong \langle C(H) \rangle$.

Proof

Let G be arbitrary. We build a new graph H which contains G as an induced subgraph via: $V_H = V_G \cup \{u, x, y, z\}$, i.e. adding 4 new vertices to G . Further, let $E_H = E_G \cup \{ux, yz\} \cup \{xv, vy \mid v \in V_G\}$,



Now $\text{ecc}(v) = 2$ for any $v \in V_G$. Nevertheless, $d(u, z) = 4$ and $d(x, z) = d(y, u) = 3$, whence $\text{ecc}(w) > 2$ for all $w \in V_H \setminus V_G$. Thus, $\text{rad}(H) = 2$ and $C(H) = V_G$, whence $\langle C(H) \rangle \cong G$.

2.9 Lemma

A graph G is isomorphic to (the graph induced by) the periphery of another graph H iff either every vertex has eccentricity 1 or no vertex does.

Proof

" \Rightarrow " We use proof by contraposition. Assume ex. $u \in V_G$ s.t. $\text{ecc}(u) = 1 < \text{diam}(G)$. In particular, $G \neq P(G)$. Now, aiming for a contradiction, assume ex. H s.t. $G \trianglelefteq H$ and $P(H) = V_G$.

As $G \neq P(G)$, we know that $H \neq G$ and $\text{diam}(H) \geq 2$.

As $u \in V_G = P(H)$, there is some $w \in V_H$ s.t. $d(u, w) = \text{diam}(H)$.

But then, $w \in P(H) = V_G$ and as $\text{ecc}(w) = 1$, we also get

$d(u, w) = 1 < \text{diam}(H)$ ∇ . Hence, $P(H)$ cannot be V_G .

" \Leftarrow " If all vertices in G have eccentricity 1 or 0, then G is complete

and $G \cong P(G)$. For the second case, assume $\text{rad}(G) > 1$.

and consider H s.t. $V_H = V_G \cup \{v\}$ contains one new vertex which is connected to everyone else, i.e. 

$E_H = E_G \cup \{vx \mid x \in V_G\}$. Then, as $\text{ecc}^G(x) \geq 2$ for all $x \in V_G$,

$$\text{ecc}(x) = \begin{cases} 2 & \text{if } x \in V_G \\ 1 & \text{if } x = v \end{cases} \quad \text{Hence, } \text{diam}(H) = 2 \text{ and}$$

$\langle P(H) \rangle = G$, as desired. 

2. Adjacency Matrices

We saw the visual benefits of studying graphs by their diagram. This is very useful to illustrate ideas and study small graphs. In applications on the other hand, when studying e.g. correlations of weather phenomena or social links, graphs tend to have thousands of vertices. Here, it is no longer practical to use neither the set- nor the diagram representation of graphs. The way computers store and analyze graphs is by using adjacency matrices.

2.10 Definition


Let G be a graph of order n with vertices $V_G = \{v_1, v_2, \dots, v_n\}$.

The adjacency matrix of G is the matrix $A_G = (a_{ij})$ of $n \times n$

defined via $a_{ij} = \begin{cases} 1 & \text{if } v_i, v_j \in E \\ 0 & \text{otherwise.} \end{cases}$ We also write $A(i,j)$ for a_{ij} .

2.11 Example

Consider G given by



. Then $A_G \in M_{4 \times 4}$

$A_G = \begin{pmatrix} 0 & 1 & 0 & 1 \\ 1 & 0 & 1 & 1 \\ 0 & 1 & 0 & 1 \\ 1 & 1 & 1 & 0 \end{pmatrix}$ is the adjacency matrix of G .

2.12 Remark

If $A_G = (a_{ij})$ is an adjacency matrix of a graph G , then

- 1) $a_{ii} = 0$ for all $1 \leq i \leq |G|$
- 2) A is symmetric.

$$3) \sum_{j=1}^{|G|} a_{ij} = \deg(v_i) \text{ and thus } \sum_{i,j=1}^{|G|} a_{ij} = \sum_{i=1}^{|G|} \deg(v_i) = 2|E|.$$

4) A_G is only unique up to reordering the vertices.

2.13 Example

Let revisit the graph $G = \begin{array}{c} v_1 \\ \bullet \\ \swarrow \quad \searrow \\ v_2 \quad v_3 \\ \downarrow \\ v_4 \end{array}$ and $A_G = \begin{pmatrix} 0 & 1 & 0 & 1 \\ 1 & 0 & 1 & 1 \\ 0 & 1 & 0 & 1 \\ 1 & 1 & 1 & 0 \end{pmatrix}$.

The fact that $A_G(2,3) \neq 0$ means that v_2 and v_3 are adjacent.

and $A(1,3)=0$ says that v_1 and v_3 are not. Now consider

$$A_G^2 = \begin{pmatrix} 0 & 1 & 0 & 1 \\ 1 & 0 & 1 & 1 \\ 0 & 1 & 0 & 1 \\ 1 & 1 & 1 & 0 \end{pmatrix} \begin{pmatrix} 0 & 1 & 0 & 1 \\ 1 & 0 & 1 & 1 \\ 0 & 1 & 0 & 1 \\ 1 & 1 & 1 & 0 \end{pmatrix} = \begin{pmatrix} 2 & 1 & 2 & 1 \\ 1 & 3 & 1 & 2 \\ 2 & 1 & 2 & 1 \\ 1 & 2 & 1 & 3 \end{pmatrix} \quad \text{Let's interpret the values of } A_G^2.$$

Now, $A_G^2(1,3) = 2$. How did we compute it?

$A_G^2(1,3) = \sum_{j=1}^4 a_{1j} a_{j3}$. Now $a_{1j} a_{j3} = 1$ iff $v_1 v_j$ and $v_j v_3$ are edges iff (v_1, v_j, v_3) is a walk of length 2 from v_1 to v_3 .

Hence $A_G^2(1,3) = \sum_{j=1}^4 a_{1j} a_{j3}$ is the number of walks from v_1 to v_3 of length 2. This generalises and provides a strong tool to study graphs.

2.14 Theorem

Let G be a graph with $V_G = \{v_1, \dots, v_n\}$ and A_G the corresponding adjacency matrix. Then the entry $A_G^k(i,j)$ is the number of possible walks from v_i to v_j of length k .

Proof

We proceed by induction on the power k . (Note that $k=0$ works, too).

$k=1$ we get that $A(i,j) = \begin{cases} 0 & \text{iff } v_i v_j \notin E_G \text{ iff there are } 0 \text{ } v_i v_j\text{-walks of length 1} \\ 1 & \text{iff } v_i v_j \in E_G \text{ iff there is 1 } v_i v_j\text{-walk of length 1.} \end{cases}$

$k \mapsto k+1$ Assume that $A^k(i,j)$ gives exactly the number of $v_i v_j$ -walks of length exactly k . Let's denote $A^k := (b_{ij})$ and $A = (a_{ij})$.

Note that there is a $v_i v_j$ -walk of length $k+1$ iff there ex.

a vertex v_e s.t. there is a $v_i v_e$ -walk of length k and an $v_e v_j$ -walk of length one. Hence

$$|\{v_i v_j\text{-walk of length } k+1\}| = \sum_{e \in \{v_i \in N(v_j)\}} |\{v_i v_e\text{-walk of length } k\}|$$

$$\stackrel{\text{l.H.}}{=} \sum_{e \in \{v_i \in N(v_j)\}} b_{ie} = \sum_{l=1}^n b_{ie} a_{ej}$$

$$= \sum_{l=1}^n A^k(i,l) \cdot A(l,j) = A^{k+1}(i,j).$$



2.15 Corollary

Let G be a graph with $V_G = \{v_1, v_2, \dots, v_n\}$ and A_G the adjacency matrix.

Then $d(v_i, v_j) = \min \{k \mid A^k(i,j) \neq 0\}$.

(Recall that $A_G^0 = I_n = \begin{pmatrix} 1 & 0 & \cdots & 0 \\ 0 & 1 & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & 1 \end{pmatrix}$).

2.16 Definition

Let G be a graph with adjacency matrix A . For every

$k \in \mathbb{N}$ we define the **skill matrix** S_k via

$$S_k = \sum_{i=0}^k A^k = I_n + A + A^2 + \dots + A^k.$$

2.17 Remark

As $S_k(i,j) = \sum_{i=0}^k A^k(i,j)$, we get that $S_k(i,j)$ is the number of $v_i v_j$ -walks of length at most k .

2.18 Example

Recall the graph $G = \begin{array}{c} v_4 \\ \backslash / \\ v_1 - v_2 - v_3 \end{array}$ with $A = \begin{pmatrix} 0 & 1 & 0 & 1 \\ 1 & 0 & 1 & 1 \\ 0 & 1 & 0 & 1 \\ 1 & 1 & 1 & 0 \end{pmatrix}$.

$$A^2 = \begin{pmatrix} 2 & 1 & 2 & 1 \\ 1 & 3 & 1 & 2 \\ 2 & 1 & 2 & 1 \\ 1 & 2 & 1 & 3 \end{pmatrix} \text{ and } A^3 = \begin{pmatrix} 2 & 5 & 2 & 5 \\ 5 & 4 & 5 & 5 \\ 2 & 5 & 2 & 5 \\ 5 & 5 & 5 & 4 \end{pmatrix}.$$

$$\text{Then } S_0 = I_4 = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}, S_1 = \begin{pmatrix} 1 & 1 & 0 & 1 \\ 1 & 1 & 1 & 1 \\ 0 & 1 & 1 & 1 \\ 1 & 1 & 1 & 1 \end{pmatrix}, S_2 = \begin{pmatrix} 3 & 2 & 2 & 3 \\ 2 & 4 & 2 & 3 \\ 2 & 2 & 3 & 2 \\ 2 & 3 & 2 & 4 \end{pmatrix} \text{ and}$$

$$S_3 = \begin{pmatrix} 5 & 7 & 4 & 8 \\ 7 & 8 & 7 & 8 \\ 4 & 7 & 5 & 7 \\ 7 & 8 & 7 & 8 \end{pmatrix}. \quad \text{This means there are for example } 4 \text{ } v_1 v_3 \text{ walks of length at most 3, namely } (v_1, v_2, v_3), (v_1, v_4, v_3), (v_1, v_2, v_4, v_3) \text{ and } (v_1, v_4, v_2, v_3).$$

INTERMEDIATE

- Consider P_4
- 1) Compute $\text{rad}(P_4)$, $\text{diam}(P_4)$, $C(P_4)$, $P(P_4)$, as well as $\text{ecc}(v_i)$ for v_1, v_2, v_3 and v_4 .
 - 2) Compute S_0, S_1, S_2 and S_3 .
 - 3) How can we read the data of 1) from the matrices on 2)?

$$A_1 = \begin{pmatrix} 0 & 1 & 0 & 0 \\ 1 & 0 & 1 & 0 \\ 0 & 1 & 0 & 1 \\ 0 & 0 & 1 & 0 \end{pmatrix} A_2 = \begin{pmatrix} 1 & 0 & 1 & 0 \\ 0 & 2 & 0 & 1 \\ 1 & 0 & 2 & 0 \\ 0 & 1 & 0 & 1 \end{pmatrix} A_3 = \begin{pmatrix} 0 & 2 & 0 & 1 \\ 2 & 0 & 3 & 0 \\ 0 & 3 & 0 & 2 \\ 1 & 0 & 2 & 0 \end{pmatrix}$$

$$S_0 = \begin{pmatrix} 1 & 1 & 0 & 0 \\ 1 & 1 & 1 & 0 \\ 0 & 1 & 1 & 1 \\ 0 & 0 & 1 & 1 \end{pmatrix} S_1 = \begin{pmatrix} 2 & 1 & 1 & 0 \\ 1 & 3 & 1 & 1 \\ 1 & 1 & 3 & 1 \\ 0 & 1 & 1 & 2 \end{pmatrix} S_2 = \begin{pmatrix} 2 & 3 & 1 & 1 \\ 3 & 3 & 4 & 1 \\ 1 & 4 & 3 & 3 \\ 1 & 1 & 3 & 2 \end{pmatrix}$$

The following theorem sums up the knowledge we acquired so far.

2.19 Theorem

Let G be a graph with $V_G = \{v_1, v_2, \dots, v_n\}$, adjacency matrix A and

skroll matrices S_k . Then the following hold.

1) $d(v_i, v_j)$ is the least k s.t. $S_k(i,j) \neq 0$.

2) $\text{ecc}(v_i)$ is the least k s.t. the i th row of S_k has no zero entries.

3) $\text{rad}(G)$ is the least k s.t. S_k contains at least one row without zero entries. or ∞ otherwise.

4) $\text{diam}(G)$ is the least k s.t. S_k does not contain any zero entries.


5) G is disconnected iff S_{n-1} contains a zero.

2.20 Definition

Let G be a graph with $V_G = \{v_1, v_2, \dots, v_n\}$. The **distance matrix** of G is the matrix $D \in M_{nn}$ s.t. $D(i,j) = d(v_i, v_j)$.

2.21 Example

Back to our example $G = \{v_1, v_2, v_3, v_4\}$. Then the



distance matrix D is $\begin{pmatrix} 0 & 1 & 2 & 1 \\ 1 & 0 & 1 & 1 \\ 2 & 1 & 0 & 1 \\ 1 & 1 & 1 & 0 \end{pmatrix}$.

2.22 Example (of a large matrix) - Erdős Number

Paul Erdős - Hungarian Mathematician, published over 1500 papers.

Consider G with $V_G = \text{all mathematicians}$, $E_G = \{(x,y) \mid x \text{ and } y \text{ published together}\}$.

Then $\deg(\text{Erdős}) > 500$ and the Erdős number of x is $d(\text{Erdős}, x)$.

Chapter 3 - TREES



The intuition for graph theoretic trees comes from actual trees in nature. Here, the stem splits into several branches that afterwards never rejoin.

3.1 Definition

A graph which does not contain cycles is called **acyclic**.


We call a graph G a **tree** if it is connected and acyclic. An arbitrary acyclic graph is called a **forest**.

In a forest, any vertex of degree 1 is called a **leaf**.


3.2 Remark

- 1) The graphs P_n , K_1 , K_2 and $K_{1,n}$ are trees for any $n \in \mathbb{N}$.
- 2) Every tree is a forest.
- 3) Every connected component in a forest is a tree.
- 4) Every subgraph of a forest is a forest.


3.3 Example




is a
tree



is a
tree.



is not
a tree.



is not a tree,
but it is a forest.

3.4 Lemma

Any tree of order at least 2 has at least two leaves.

Proof

Let T be a tree with $|T| \geq 2$. (In particular, T is connected).

Consider a path of maximal length $p = (v_0, v_1, \dots, v_n)$ in T . As $|T| \geq 2$,

we know that $v_0 \neq v_n$. We claim that v_0 and v_n are leaves,

i.e. $\deg(v_0) = \deg(v_n) = 1$. We execute the argument for v_0 .

As usual, we know that $N(v_0) \subseteq \{v_1, v_2, \dots, v_n\}$. Let $u \in N(v_0)$

arbitrary, i.e. $u = v_i$ for some $i \geq 1$. But then $(v_0, v_1, \dots, v_{i-1}, v_i, v_0)$

is a closed walk which is a cycle for all $i \geq 2$. As T does not contain cycles, we conclude that $i = 1$ and v_1 is the only neighbour of v_0 . Hence $\deg(v_0) = 1$ and v_0 is a leaf.

The argument for v_n is analogous. \square


3.5 Definition (Tree Pruning)

Let T be a tree of order at least 3. We denote by

T^- the induced subgraph of T obtained by deleting all leaves of T .

3.6 Example

i) If $T = P_7$ the path of length 6, then $T^- = P_7^- = P_5$ is the path of length 4:



ii) If $T = K_{1,n}$ the complete bipartite graph, then $T^- = K_1 = E$, consists of one vertex only:



3.7 Lemma

Any tree T of order n has exactly $n-1$ edges.

Proof

We proceed by induction on $|T|$.

If $|T|=1$, then $T=K_1$, which has zero edges and the claim holds.

Now assume we know that any tree of order n has exactly $n-1$ many edges and consider T of order $n+1$ arbitrary.

By Lemma 3.4, T has a leaf u . Clearly, $T-u$ is still connected and of order n , whence $T-u$ has exactly $n-1$ edges. As u was a leaf in T , T has exactly one edge more than $T-u$, whence $|T|=n=(n+1)-1$, as desired. \square

3.8 Corollary

A forest of order n , consisting of k -many connected components, has exactly $n-k$ many edges.

We will now see that given a graph G is connected, Lemma 3.5 is not only a necessary, but even a sufficient condition for G to be a tree.

3.9 Theorem

A graph G of order n is a tree iff it is connected and has exactly $n-1$ many edges.

Proof

" \Rightarrow " Clear by definition of a tree and Lemma 3.5.

" \Leftarrow " Assume G is connected of order n and contains exactly $n-1$ many edges. If G contains a cycle, take any edge e , within the cycle and consider $G-e$. Then $G-e$ is still connected and of order n . If $G-e$ still contains a cycle, we proceed likewise and after $k \leq n-1$ many steps we obtain a graph $G-\{e_1, e_2, \dots, e_k\}$ which is of order n , connected and without cycles, whence it is a tree. But $G-\{e_1, e_2, \dots, e_k\}$ has $(n-1)-k < n-1$ many edges, contradicting Lemma 3.5. \square

3.10 Theorem

A graph of order n is a tree iff it is acyclic and has $n-1$ many edges.

Proof

" \Rightarrow " clear.

" \Leftarrow " Assume G is of order n with $n-1$ many edges and acyclic, i.e. G is a forest. But by Corollary 3.6., if G has k -many connected components than $|G| = n-k = n-1$, whence $k=1$ and G is connected and hence a tree. \square

3.11 Corollary | Summary

Let G be a graph of order n . Then tfae:

- 1) G is connected and acyclic (i.e a tree)
- 2) G is connected and has $n-1$ many edges.
- 3) G is acyclic and has $n-1$ many edges.

3.12 Home work

Every edge in a tree is a bridge.

3.13 Lemma

For any two vertices $u, v \in V_T$ in a tree T , there is a unique uv -path.

Proof

As T is connected, there clearly is a uv -path for any $u, v \in V_T$.

Now assume that $p_1 = (u = x_0, x_1, \dots, x_e = v)$ and $p_2 = (u = y_0, y_1, \dots, y_e = v)$

are two distinct uv -paths. Then $\langle p_1 \cup p_2 \rangle$ is again a tree.

Let i be minimal s.t. $x_i \neq y_i$. Then $\langle p_1 \cup p_2 \rangle - y_i y_{i+1}$ is still connected, contradicting the fact that every edge in a tree is a bridge.

◻

3.14 Corollary

Let T be a tree and $v \in V_T$. Then $\text{ecc}(v)$ is the length of the longest path starting from v .

3.15 Lemma

Let T be a tree of order at least 2. Consider $u, v \in V_T$ s.t. $\text{ecc}(v) = d(u, v)$. Then u is a leaf.

Proof

Let $p = (v = x_0, x_1, \dots, x_e = u)$ be the unique vu path. If u were not a leaf, then it had at least one neighbour $w \notin p$.

But then $(v = x_0, x_1, \dots, x_e, w)$ would be a path starting in v and longer than p , contradicting Corollary 3.12. ◻

3.16 lemma

Let T be a tree of order at least 3. Then $C(T) = C(T^-)$.

Proof

1) Show that $C(T) \subseteq T^-$, i.e. $C(T)$ contains no leaf.

To this end, let u be a leaf and v its unique neighbour.

As $|T| \geq 3$, v is not a leaf itself and $d(u,w) = d(v,w) + 1$ for any $w \in V_T \setminus \{u\}$, whence $\text{ecc}(u) > \text{ecc}(v)$ and hence $u \notin C(T)$.

2) Show that $\text{ecc}^{T^-}(v) = \text{ecc}^T(v) - 1$ for every nonleaf $v \in V_T$.

To that end, consider an arbitrary non-leaf $v \in V_T$ and pick $u \in V_T$ s.t. $d(v,u) = \text{ecc}(v)$. By 3.13, u is a leaf. Let

p be the unique vu -path in T and note that u is the only leaf on p . Hence only u will be deleted from p in T^- . As this holds for all paths in T starting in v of length $\text{ecc}(v)$, we obtain that $\text{ecc}^{T^-}(v) = \text{ecc}^T(v) - 1$, as desired.

3) We conclude from 1) + 2) that for any vertex $v \in T^-$,

$\text{ecc}^{T^-}(v) = \text{ecc}^T(v) - 1$, whence $v \in C(T)$ iff $v \in C(T^-)$ (and $\text{rad}(T^-) = \text{rad}(T) - 1$).

3.17 Lemma

Let T be a tree. Then $C(T)$ is either K_1 or K_2 .

Proof

We do induction on $|T|$. If $|T|=1$, then $T=K_1$ is its own center and we are done. Similarly for $|T|=2$, where $T=K_2$.

Now assume that the claim holds for all trees of order $n \geq 3$, and consider a tree T with $|T| = n+1$ arbitrary.

By 3.16, we know that $C(T) = C(T')$. By 3.4 we know that T contains at least two leaves, whence $|T|-1 \leq |T|-2 < n$. Hence, by I.H., $C(T) = C(T')$ is either K_2 or U_3 as desired. 

3.18 Lemma

Let T be a tree of order n and G an arbitrary graph s.t. $\delta(G) \geq n-1$. Then G contains T as a subgraph.

Proof

We use induction on $|T|$.

If $|T|=1$, then $T=K_1$ is a subgraph of any graph G .

Now assume we proved the claim for all trees of order at most n .

Consider T with $|T|=n+1$ and G with $\delta(G) \geq n$ arbitrary.

Let u be a leaf of T and denote by $T' := T - u$. Then $|T'|=n$ whence T' can be seen as a subgraph of G . Let v be the unique neighbour of u in T . Then $\deg^G(v) \geq \delta(G) \geq n$, but as

$|T'|=n$ and v cannot be its own neighbour, there exist some $u' \in G$ adjacent to v and not contained in T' . Hence, the subgraph $(V_{T'} \cup \{u'\}, E_{T'} \cup \{v, u'\})$ is the desired subgraph of G isomorphic to T . 

Summary

- 1) A tree of order n contains exactly $n-1$ edges.
- 2) Any tree of order at least two contains at least two leaves.
- 3) A graph of order n is a tree iff it is connected of size $n-1$.
- 4) A graph of order n is a tree iff it is acyclic and of size $n-1$.
- 5) A graph is a tree iff for any vertices u, v there is a unique uv -path.
- 6) The centre of any tree is either K_1 or K_2 .
- 7) Any graph G contains any tree of order at most $\delta(G)+1$ as a subgraph.

Spanning Trees

3.19 Definition

Let G be any graph. We call a subgraph $T \subseteq G$ a **spanning tree** for G if it is a tree and contains all vertices of T .


3.20 Remark

From the previous chapter it is clear that a spanning tree of a graph G of order n has n many vertices and $n-1$ many edges.


3.21 Examples

Consider the following graphs and spanning trees.

1) $G = C_6$, a possible spanning tree:



2) $G = K_5$, possible spanning trees:



3.22 Lemma

Every connected graph contains at least one spanning tree.

Proof


Assume G is connected and let T be a subgraph of G of maximal order s.t. T is a tree. We need to show that $V_T = V_G$. Otherwise, as G is connected, there is some vertex $u \in V_G \setminus V_T$ which is adjacent to some vertex $v \in V_T$. Now, consider the new subgraph $\tilde{T} = (V_T \cup \{u\}, E_T \cup \{uv\})$. Its $\deg(\tilde{T})(u) = 1$, u is not contained in any cycles in \tilde{T} , whence \tilde{T} is still a tree. As this contradicts maximality of $|T|$, we conclude that T must contain all vertices of G , whence it is a spanning tree for T .

3.24 Definition

A function $w: E_G \rightarrow \mathbb{R}$ is called a **weight function** on G .
 A graph G together with a weight function, i.e. the triple (V_G, E_G, w) is called a **weighted graph**.

3.25 Example - Visualisation


We visualise the weighting of a graph by denoting the weight $w(e)$ on top of the edge e , e.g.





Here the weight function of an edge $e=uv$ is given by the (birds eye) distance between u and v .

3.26 Example

Consider the following weighted graph.



We can find several spanning trees. Let's name some and compute their weight.



3.27 Definition

Let (G, w) be a connected weighted tree. A minimum-weight spanning tree T is a spanning tree of G s.t. the sum of the weights of its edges is minimal among all possible spanning trees of G , i.e. if T' is another spanning tree then $\sum_{e \in E_T} w(e) \leq \sum_{e \in E_{T'}} w(e)$.

Now how can we find a minimal spanning tree effectively?

Consider the following algorithm:

3.28 Kruskal's Algorithm (1956)

Consider the set of vertices as a forest $F = (V_S, \phi)$ where

each vertex is a maximal subtree of F . Let $E := E_{\mathcal{S}}$.

While (F is not a free $\mathbf{1} \in \mathcal{E} \neq \emptyset$)

Pick $e \in E$ of minimal weight. Let $E := E \setminus \{e\}$.

If e connects two trees in F , let $E_F = E_F \cup \{e\}$.
 (i.e. $F \cup e$ is still acyclic)

This algorithm stops after at most $|E_S|$ many repetitions.





329 Example

We apply the algorithm on


Let us mark edges we

add to F green and the ones


We disregard, red.




4)



5)




6)




$$E := E - \{e,f\}$$

$$E_F := E_F \cup \{e,f\}$$


7)



8)




9)



$$E := E - \{d,e,f\}$$

$$E_F := E_F$$



$$E := E - \{d,f,g\}$$

$$E_F := E_F \cup \{d,f,g\}$$

Here the algorithm stops, as

$$F = (V_G, E_F) \text{ with } E_F = \{ab, ac, gh, eg, dg, cd, df\}$$



is a single tree whence the conditions in

the while loop are violated.

(The output is the spanning tree F. Note that the second condition in the while loop was still valid, as $E = \{ad, de\} \neq \emptyset$).

INTERMEZZO

Use Kruskal's algorithm to find a minimal spanning tree for each of the two given graphs. Is this spanning tree unique? Argue your answer.



3.30 Theorem

Kruskal's algorithm is correct, i.e. it always terminates and its output is a minimum-weight spanning tree.

Proof

1) Termination: As after $\|G\|$ -many steps the condition $E \neq \emptyset$ is violated, the algorithm always terminates.

2) The output F is a spanning tree:

As $V_F = V_G$, it clearly contains all vertices of G .

Further, in each step the regarded edge e either connects two disconnected trees into one larger tree, or, if it would connect two vertices of the same subtree in F , is disregarded. Hence after each step, F is still a forest, i.e. acyclic.

It remains to show that F is connected. If the algorithm stops because F is a tree, then it is clearly connected. If it stops, because we went through all the edges, then any edge of G not contained in F would connect two vertices of the same connected component. Thus F has as many connected components as G , which is one, as G is connected.

3) F is a minimum-weight spanning tree.

Aiming for a contradiction, assume this is not the case.

Let $\{e_1, \dots, e_m\}$ be all the edges in F , enumerated in the order they were added to F by the algorithm.

Among all possible minimum-weight spanning trees, let T be one that agrees with F on the largest initial segment of $\{e_1, e_2, \dots, e_{k-1}\}$, i.e. if k is the smallest index s.t. $e_{k+1} \notin T$, then there is no minimum-weight spanning tree which contains $\{e_1, e_2, \dots, e_k, e_{k+1}\}$.

As by assumption F is not minimum-weight, we have $k < n-1$.

As T is a spanning tree which does not contain e_{k+1} , we know that $T + e_{k+1}$ contains a cycle. C. As F did not contain cycles, there is one edge $e \in C \setminus T$ which is not in F .

Now $T + e_{k+1} - e$ is a connected graph of order n and size $n-1$, whence still a spanning tree. It contains the edges $\{e_1, \dots, e_k, e_{k+1}\}$, hence it can no longer be of minimum weight. This means that $w(e_{k+1}) > w(e)$.

But as $e \notin F$ and in particular $e \notin \{e_1, \dots, e_k\}$ this means e was available at the step of the algorithm after we added e_k and of less weight than e_{k+1} . This contradicts the assumption that the algorithm chooses the edge of minimal weight which keeps F acyclic. \square

3.31 Lemma

If G is a connected weighted graph s.t. distinct edges have distinct weights, then there is a unique minimum-weight spanning tree.

Proof

Homework, will be included after.

Chapter 4 - Euler and Hamilton

Imagine

A salesperson with their wagon wants to pass by every street in his neighbourhood to sell their goods. Of course, they want to minimize efforts, so they would like to avoid passing the same street twice. These type of problems are considered when discussing Eulerian graphs.

Then, as only few people buy, they switch to their car and only visit a central place in each city of the area. Again, to improve efficiency, they only want to visit each city once. This type of problem is studied when discussing Hamiltonian graphs.


How do these two problems differ? Let's find out!

4.1. Definition

We call a trail on a graph G an Eulerian trail if it contains every edge of G . We call it an Eulerian circuit if it is a closed Eulerian trail. Finally, the graph G itself is called an Eulerian graph iff it contains an Eulerian circuit.


4.2 Examples

1)  is not Eulerian but has Eulerian trail.

2)  is indeed Eulerian.

3) Any cycle C_n is clearly Eulerian.

4) Any path of length $n \geq 1$ is not Eulerian but has an Eulerian trail.

5)  is indeed Eulerian, but

6)  does not even contain an Eulerian trail.

7) Generally, every K_{2n+1} is Eulerian and every K_{2n+2} does not even contain an Eulerian trail. for $n \geq 1$.


4.3 Observation

Consider the graph K_5 . We observe the following properties:

1) K_5 is Eulerian:


(a,b,c,d,e,a,d,b,e,c,a)

is an Eulerian circuit.




2) Every vertex of

K_5 has an even degree.



3) We can partition E_K

into cycles (i.e. find mutually edge-disjoint cycles that together use all edges of K_5).



These three properties do not appear together by coincidence. It turns out, they are equivalent to each other.

4.4. Auxiliary Lemma

Let G be a connected graph with $|G| \geq 2$.

If $\deg(v)$ is even for all $v \in V_G$, then G contains a cycle C . Moreover, $G - C$ still contains a cycle or is $E_{|V_G|}$.

Proof

Assume G is as above. If G would not contain a cycle, it was a tree. But then it had to contain a leaf v .

But then $\deg(v) = 1$ is not even. For the "moreover"-

Part observe that $\deg^{G-C}(v) = \begin{cases} \deg^G(v) - 2 & \text{if } v \in V_C \\ \deg^G(v) & \text{else} \end{cases}$, hence

still even. Then each connected component of $G - C$ still contains a cycle (whence so does $G - C$), or is of order 1. \square

4.5 Theorem (Euler-Hierholzer-Vetlesen)

Let G be a connected graph. The following are equivalent:


- 1) G is Eulerian.
- 2) Every vertex of G is of even degree.
- 3) The edge set of G can be partitioned into a set of edge-disjoint cycles.

4.6 Corollary

A graph contains an Eulerian trail iff either each vertex has even degree or there are exactly two vertices of odd degree.

INTERMEZZO

- 1) Consider the following graph. Is it Eulerian? If so, argue your answer by



- a) Giving an Eulerian circuit.
- b) using the degree sequence.
- c) Partitioning the edges into cycles.

- 2) For which n and m is $K_{n,m}$ Eulerian?

Proof of Theorem 4.5

As all three clearly hold for $|G|=1$, we may assume that $|G|>1$.

1 \Rightarrow 2) Assume G is Eulerian. Let Q be an Eulerian circuit of G . Now consider $v \in V_G$ arbitrary. Without loss of generalisation, we may assume that Q does not start with v . Now, every appearance of v on Q corresponds to two distinct edges involving v , the one leading into v and the one leading away from v . As Q is Eulerian, it uses all edges incident with v . Whence in total there is an even number of edges incident with v and $\deg(v)$ is even. ✓

2 \Rightarrow 3) Assume G only contains vertices of even degree.

By Lemma 4.4, G contains at least one cycle C . We proceed by induction on the number n of cycles in C .

$n=1$: If G contains only one cycle, then $G=C_{|G|}$ and hence the desired partition of edges is just the cycle G itself.

$n \Rightarrow n+1$: Now assume every graph containing at most n -many cycles allows a partition into edge-disjoint cycles. Consider any connected G with $(n+1)$ -many cycles. Pick an arbitrary cycle C in G . Then, as in 4.4., in $H := (V_G, E_G - E_C)$, every vertex still has even degree. Now, every connected component of H contains at most n -many cycles and

By induction hypothesis, we can partition each connected component of H , and hence H itself, into edge-disjoint cycles. Once we add C to this partition, we obtain the desired partition of G . ✓ (Note that this gives you a cooking recipe of how to find cycles).

3 \Rightarrow 1: Assume the edge set of G can be partitioned into k -many sets S_1, S_2, \dots, S_k s.t. the edges of each S_i form a cycle. Let Q be a circuit of maximal length in G s.t. the edges of Q equals the union of some sets S_i , i.e. such that there is $I \subseteq \{1, \dots, k\}$ with $E_Q = \bigcup_{i \in I} S_i$.

As the S_i are pairwise disjoint, we know that Q contains either no edge from S_i or all edges from S_i for ever $i \in I$.

Now if $E_Q = E_G$, then G is Eulerian and we are done.

Otherwise, there is some edge not contained in Q , but incident with a vertex v in Q . The edge must be contained in exactly one S_l with $l \notin I$. Note that Q and S_l have no common edges, but they share the vertex v . Hence we may glue the circuit Q and the cycle S_l at v and obtain a new circuit Q' longer than Q with $E_{Q'} = \bigcup_{i \in I \cup \{l\}} S_i$, contradicting our choice of Q .

Hence Q contained all edges of G and hence G is Eulerian. ✓



4.2 - Hamilton

Sir William Rowan Hamilton (1805-1865)

- Irish pure mathematician
- contributions to optics, mechanics and algebra.
- also invented a game (The Icosian Game) build on graph theory (bought by Jaques and Son, huge failure)

4.7 Definition

Let G be a graph. A **Hamiltonian path** is a path in G which uses all vertices of G . A **Hamiltonian cycle** is a cycle in G which uses all of V_G .

We call G **traceable** if it contains a Hamiltonian path and we call it **Hamiltonian** if it contains a Hamiltonian cycle.


4.8 Remark

- 1) Every Hamiltonian graph is traceable but not vice versa.
- 2) Traceable graphs are connected.
- 3) If $|G|=n$, then G is Hamiltonian iff it contains C_n as a subgraph and it is traceable iff it contains P_n as a subgraph.

4.9 Examples


1) C_6 is Hamiltonian via

All vertices have even degree.




2) K_4 is Hamiltonian via

All vertices have odd degree.



3) The graph G_1 is Hamiltonian.

There are vertices of even and odd degree.




4) The graph G_2 is not Hamiltonian.

Every vertex has even degree.




5) The graph $K_{1,3}$ is not Hamiltonian.

All vertices have odd degree.




6) The path P_4 is not Hamiltonian.

There are vertices of even and odd degree.



INTERMEZZO



1) Play the Icosian game!

Place the numbers 1-20 into


the vertices of the graph on the left

st. $(1, 20)$ as well as $(k, k+1)$ are edges

in the graph for all $k \in \{1, \dots, 19\}$

2) Is the graph Hamiltonian? Is it Eulerian?

Solution and name-motivation for the icosian game:



(gr. ΕΥΚΟΣΙ ("ikosi") = 20 \Rightarrow 20 vertices)

played on a dodecahedron, δοδεκά - gr. dodeca, twelve, εδρα ("edra") = seat, face

\hookrightarrow polyhedron with 12 faces, 20 vertices.

4.10 Remark

While it is rather easy to decide whether a graph is Eulerian (P-TIME, $O(|G|^2)$), it is surprisingly hard to do the same for Hamiltonian graphs. This problem is known to be NP-complete and still we did not manage to find an equivalent condition for Hamiltonianity (other than containing $C_{|G|}$ as a subgraph, which is basically the definition).

We hence see, even though the Eulerian graph problem and the Hamiltonian graph problem seem so similar, their resolution requires very different levels of efforts.

The best we can do at the moment is give some sufficient criteria.

4.11 Theorem (Dirac)

Let G be s.t. $|G| \geq 3$. If $\delta(G) \geq \frac{n}{2}$, then G is Hamiltonian.

Proof

Consider G arbitrary s.t. $|G|=n \geq 3$ and $\delta(G) \geq \frac{n}{2}$.

Then G is necessarily connected (think why).

Consider a path $p = (v_1, v_2, \dots, v_k, v_k)$ of maximal length in G .

We claim that there is some $j \leq k$ s.t. $v_j \in N(v_k)$ and

$v_{j+1} \in N(v_1)$, i.e.



is a subgraph

of p . Note that as usual, as p is of maximal length,

all neighbours of v_1 and v_k must be on p . As $\delta(G) \geq \frac{n}{2}$,

v_k has at least $\frac{n}{2}$ many neighbours v_j in p . Aiming for

a contradiction, assume for every neighbour $v_j \in N(v_k)$,

$v_{j+1} \notin N(v_1)$: Then there are at least $\frac{n}{2}$ many vertices in p

which are not neighbours of v_1 . This now yields

the desired contradiction, as all neighbours of v_1 are on p


and thus $\deg(v_1) \leq (k-1) - \frac{n}{2} \leq (n-1) - \frac{n}{2} = \frac{n}{2} - 1 < \frac{n}{2}$,

contradicting $\delta(G) \geq \frac{n}{2}$ (so in part. $\deg(v_1) \geq \frac{n}{2}$). \square

Hence there is some j , s.t. $v_1 v_{j+1}$ and $v_j v_k$ are edges,

which leads to the existence of a cycle

$C = (v_1, v_2, \dots, v_{j-1}, v_j, v_k, v_{k-1}, \dots, v_{j+1}, v_i)$, i.e.



Finally, we claim that C is indeed a Hamiltonian cycle, i.e.

it contains all vertices of G . Otherwise, as G is connected, there is a vertex u in $G \setminus C$ which is adjacent to one vertex v_i in C . But as C is a cycle, we can form a new path starting $(uv_i \dots)$ and then travelling through all $k-1$ many vertices of C . This path is longer than P , contradicting our choice of P .

Hence, C indeed contains all vertices of G whence it is a Hamiltonian cycle and G is Hamiltonian.

This concludes the proof to our first sufficient criterion on Hamiltonicity. The following fact is a strengthening of Dirac's theorem and can be proven analogously.

4.12 Fact (Ore, 1960)

Let G be a graph of order $n \geq 3$. Suppose for every pair of non-adjacent vertices u, v we have that $\deg(u) + \deg(v) \geq n$. Then G is Hamiltonian.

Note that now Dirac's Theorem is a mere corollary of Ore's theorem.

We want to achieve yet another sufficient criterion for Hamiltonicity. This leads us to the so-called independence number.

4.13 Definition

Let G be a graph. A set $S \subseteq V_G$ of vertices is called an independent set if any two vertices in S are nonadjacent.

The independence number $\alpha(G)$ of G is the maximal size of an independent set.

4.14 Examples

• $\alpha(E_n) = n$, as V_{E_n} is an independent set.


• $\alpha(K_n) = 1$, as any two vertices are adjacent.

Actually, the converse also holds, i.e. $\alpha(G) = 1$ iff G is complete.

• $\alpha(K_{n,m}) = \max\{n, m\}$, as any set is independent iff it is contained in one of the parts.



Consider the given graph
(Petersen graph). Find $\alpha(G)$
and $K(G)$. What type of graph
is G ? Is G Hamiltonian?




4.15 Notation

If p is a path and x, y are two vertices on p , then we denote by $p[x,y]$ the subpath on p from x to y , i.e. for $v_1 \dots v_2 \dots v_3 \dots v_4 \dots v_5 \dots v_6 \dots v_7$

$$p[v_6, v_3] = (v_6, v_5, v_4, v_3).$$

Similarly, if C is a cycle and $x, y \in C$, $x \neq y$, then we denote by $C^+[x,y]$ the xy -path on C in clockwise direction and by $C^-[x,y]$ the xy -path on C in counter clockwise direction.

e.g. if C is , then $C^+[x,y] = (v_1, v_2, v_3, v_4)$ and $C^-[x,y] = (v_1, v_8, v_7, v_6, v_5, v_4).$

Finally, for sequences $s = (x_1, \dots, x_e)$, $t = (y_1, \dots, y_t)$ we define

$s \cdot t := (x_1, \dots, x_e, y_1, \dots, y_t)$ to be the concatenation of both.

4.16 Theorem (Chvátal, Erdős, 1972)

Let G be a graph of order at least 3.

If $\chi(G) \geq \alpha(G)$, then G is Hamiltonian.

Proof

Let G be as above, i.e. $|G| \geq 3$, $\chi(G) \geq 1$, $\chi(G) \geq \alpha(G)$.

- First we argue that $\chi(G) \geq 2$. Otherwise $\chi(G) = \alpha(G) = 1$, whence

G is a complete graph. As further $\chi(K_n) = n - 1$, G would be K_2 , contradicting $|G| \geq 3$.

- Hence now we know that $\chi(G) \geq 2$. By 1.41(8), we know that $\delta(G) \geq \chi(G) \geq 2$, whence by 1.39, G contains a cycle.


Now consider a cycle C of maximal length in G . We claim that C is Hamiltonian.

Aiming for a contradiction, assume C is not Hamiltonian,
i.e. there is some vertex $v \notin C$.

Let H be the connected component of v in $G \setminus C$.


i.e.

$$G =$$



Now, we list all elements of C which are connected to some vertex in H in clockwise order:

$$\{c_1, c_2, \dots, c_r\} \quad (\text{s.t. } c_j \in C^+ [c_{j-1}, c_{j+1}]), \text{ i.e.}$$



where each c_i is adjacent to some $h_i \in H$.


Claim 1: No two c_i 's are consecutive vertices in C .

Proof: Otherwise assume there is an i s.t. c_{i+1} is the clockwise successor of c_i . Let p be a path from h_i to h_{i+1} in H .

Then $C^+ [c_{i+1}, c_i] \setminus \{c_i, h_i\} \cup p \cup h_i f_{i+1}$ is a cycle strictly longer than C , contradicting our assumptions.

i.e.

$$G =$$




$$|C_2| > |C| \quad \text{.}$$

Now that no two c_i and c_j are clockwise successors, we can define the set $D = \{d_1, d_2, \dots, d_r\}$ where each d_i is the clockwise successor of c_i in C and we get that

$$\{c_1, c_2, \dots, c_r\} \cap \{d_1, d_2, \dots, d_r\} = \emptyset.$$

So we obtain




Note that no d_i is adjacent to any vertex in H , as otherwise we get that $d_i = c_j$ for some j . Hence also $d_i \notin H$ for any d_i .

Claim 2: $\{c_1, c_2, \dots, c_r\}$ is a cut set for G .

Proof: This is clear as any path from v to a vertex in C has to pass through one of the vertices in $\{c_1, \dots, c_r\}$, so $G - \{c_1, \dots, c_r\}$ is disconnected.

Consequently, as $\kappa(G)$ is the size of a smallest cut set, we obtain that $r \geq \kappa(G) \geq 2$.


Claim 3: There are d_i and d_j which are adjacent, i.e.



Proof: Consider the set $X := \{d_1, d_2, \dots, d_r, v\}$, and recall that there is no edge between any d_i and v . It's $|X| = r+1 \geq \kappa(G)+1 > \alpha(G)$, X cannot be an independent set, whence at least one pair d_i, d_j must be adjacent.

Now we are ready for our final contradiction: We produce a cycle ℓ longer than C . Assume $d_i d_j$ is an edge and $i < j$. Let $g_{hi,hj}$ be a path in H from h_i to h_j .

Now define $\hat{C} = (c_i) \cup q_{hi, hi} \cup \bar{C}[c_j, d_i] \cup C^+[d_j, c_i]$, i.e.



Note that \hat{C} uses all edges of C except the edges $c_i d_i$ and $c_j d_j$. Instead it uses at least the three additional edges $c_i h_i$, $h_j c_j$ and $d_i d_j$. Hence, \hat{C} is strictly longer than the cycle C , which contradicts our choice of C .
Conclusively, C must contain all vertices of G and hence is Hamiltonian. □

So far, we have encountered sufficient criteria for Hamiltonian graphs using the degree of vertices and the independence number. We will conclude the chapter by providing a last sufficient criterium using a new concept - forbidden subgraphs.

4.17 Definition

Let H and G be graphs. We say that G is H -free if H is not (isomorphic to) an induced subgraph of G .

Moreover, if S is a collection of graphs, then we call G S -free iff G is H -free for any H in S .

4.18 Example


The Petersen graph

is indeed C_3 -free

is not C_5 -free

is not E_4 -free

is indeed E_5 -free



Recall:


$$\kappa(G) = 3$$

$$\alpha(G) = 4$$


G is hence also $\{C_3, E_5\}$ -free.

4.19 Notations

Let Z_1 be the graph $Z_1 =$



and $N =$



Further, we call the graph $K_{1,3}$ the claw,

based on its shape: $K_{1,3} =$




4.20 Theorem (Goodman, Hedetniemi, '74)

Let G be 2-connected and $\{K_{1,3}, Z_1\}$ -free, then G is Hamiltonian.

Proof

As $\delta(G) \geq \kappa(G) \geq 2$, we get that G contains a cycle. Consider such a cycle C of maximal length. We claim that C is Hamiltonian.

Otherwise, as G is connected, there was a vertex $v \in V_G$ not on C , but adjacent to some vertex x on C , i.e.




Denote by y and z the neighbours of x on C .

Note that yz is not an edge as otherwise


replacing the subsequence (y, x) in C by (y, v, x)

would yield a cycle longer than C. Similarly, v₂ is not an edge. Consequently, the induced subgraph on S = {x, y, z, v₂} is either



$$= K_{1,3}$$

or



$$= Z_3$$

both of which contradict our assumptions on G. \square

The following final condition is now easy to verify:

4.21 (Duffus, Gould, Jacobson, 1980)

Let G be a $\{K_{1,3}, N\}$ -free graph.

- i) If G is connected, it is traceable.
- ii) If G is 2-connected, it is Hamiltonian.

Note that neither of these are necessary for G to be Hamiltonian. Indeed, for any graph H there is a Hamiltonian graph G which contains H as an induced subgraph.

Chapter 5 - Planarity


(lat. planaris - flat, level)


5.1 Definition

A graph G is called **Planar** if it can be drawn on a plane s.t. its edges at most intersect in their end vertices. Any such drawing is then called a **planar representation** or a **planar embedding**. If G is not planar, it is called **nonplanar**.


5.2 Examples

1) Clearly, trees, cycles and empty graphs are planar.

2) The graph  is planar and this is a planar representation.

3) The graph $K_{2,3}$  is planar, but this is not a planar representation.

One planar representation is given by



In order to prove that a graph is planar, we "just" have to provide one planar representation. But these can be very hard to find.

In order to prove that a graph is not planar, we would have to check "all" possible representations. This is not feasible.

But we can help ourselves by throwing some math on the problem.

To this end, we need some more terminology.


5.3 Definition

A region in a planar rep. is a maximal area of the plane s.t. any two points within can be connected by a curve which does not intersect or touch any part of the graph.

Regions which are completely bounded (=surrounded) by graph edges are called interior regions. The unique non-interior region is called exterior region.

5.4 Example


Consider the planar representation




We can find the regions by

considering this representation as a cookie cutter which we use to part the dough on our table surface.

So the regions are:



=>




where R₁-R₄ are interior and R₅ is the exterior region.

5.5 Definition

Let Γ be a planar representation of G in \mathbb{R}^2 . We say that an edge e is incident with a region R if for every point x on the edge e and $\epsilon > 0$ there is a point y in \mathbb{R} st. $d(x, y) < \epsilon$ (or $B_\epsilon(x) \cap R \neq \emptyset$). We say that e is a bound for R , if it is incident with R and at least one other region. We denote the number of bounds for a given region R by $b(R)$ and call it its bound degree while $B(R) = \{e \in E(G) \mid e \text{ is a bound for } R\}$ is called the boundary of R .

5.6 Example

Consider G given by its planar representation



Then e_6 is a bound of R_2 , as it is incident with R_1 and R_2 , but e_7 is not a bound of R_2 , as it is only incident with R_2 .
further, $B(R_2) = \{e_1, e_2, e_3, e_4, e_5, e_6\}$

and $b(R_2) = 6$. Also, $B(R_3) = \{e_1, e_2, e_3\}$ and $B(R_1) = \{e_4, e_5, e_6\}$.

5.7 Remarks

- It very proper treatment needs a good fusion of real analysis with topology and exceeds our time frame (nice thesis topic)

- Any edge is incident with either 1 or 2 regions, whence

$$\sum_{R \text{ region}} b(R) = \sum_{R \text{ region}} |B(R)| = 2|\{e \mid e \text{ is a bound for some } R\}| \leq 2|E|$$

- Any region either has no bounds or at least three.

- A region has no bound iff it is the only region and G is a forest.

5.9 Fact

Assume Γ is the planar representation of a graph G with regions R_1, R_2, \dots, R_r . Then

1) Either G is a forest and $r=1$ and $b(R_i)=0$ or


2) G contains a cycle and $r \geq 2$ with

$$3r \leq \sum_{i=1}^r b(R_i) \leq 2|E|.$$


INTERMEZZO

For any of the graphs below, let $n:=|G|$, $m=|G|$ and r be the number of regions. Compute the data. Investigate the triple (n, m, r) .


Do you notice anything?




$$\begin{aligned} n &= 4 \\ m &= 4 \\ r &= 2 \\ \sum_{i=1}^r b(R_i) &= 8 = 2m \end{aligned}$$




$$\begin{aligned} n &= 5 \\ m &= 10 \\ r &= 4 \\ \sum_{i=1}^r b(R_i) &= 14 = 2m \end{aligned}$$




$$\begin{aligned} n &= 6 \\ m &= 15 \\ r &= 5 \\ \sum_{i=1}^r b(R_i) &= 20 = 2m \end{aligned}$$



$$\begin{aligned} n &= 7 \\ m &= 21 \\ r &= 6 \\ \sum_{i=1}^r b(R_i) &= 12 < 2m \end{aligned}$$



$$\begin{aligned} n &= 8 \\ m &= 28 \\ r &= 7 \\ \sum_{i=1}^r b(R_i) &= 24 = 2m \end{aligned}$$



$$\begin{aligned} n &= 9 \\ m &= 36 \\ r &= 8 \\ \sum_{i=1}^r b(R_i) &= 0 < 2m \end{aligned}$$

Note that $r=m-n+2$ for all graphs. Also $\sum b(R_i)=2m$ iff every edge is part of a cycle.

5.10 Observation

Let e be the edge of a planar graph G . Then e is a bound for some region R if e is part of a cycle in G .

We now will prove that the relation discovered in the Intermezzo holds for any planar representation.

5.11 Theorem (Euler's Formula)

Let Π be any planar representation of a connected graph G .

Then for $|G|=n$, $\|G\|=m$ and r the number of regions in Π , we have.

$$n-m+r=2.$$

5.12 Corollaries

1) As consequently $r = \|G\| - |G| + 2$, we get that for a planar graph G , the number of regions is independent from the chosen planar representation. We can hence say that G has r -many regions.

2) If G is a planar graph on k -many connected components C_1, \dots, C_k then each C_i is planar. If C_i has r_i -many regions, note that G has $\sum_{i=1}^k r_i - (k-1)$ many regions as all components share the common exterior region in a joint embedding. Thus the number of regions r of G is

$$\begin{aligned} r &= (\sum r_i) - (k-1) = \sum ((\|C_i\| - |C_i| + 2) - (k-1)) \\ &= \|G\| - |G| + 2k - k + 1 = \|G\| - |G| + k + 1. \end{aligned}$$

Hence, for arbitrary planar G with $|G|=n$, $\|G\|=m$ and $r(G)=r$,

we get

$$n-m+r=k+1$$

Proof of Theorem 5.11

Let G be a connected planar graph in any given planar representation with regions R_1, R_2, \dots, R_r . Let $|G|=n$.

We prove the theorem by induction on $\|G\|=m$

$m=0$: If G has no edges, then $G=K_1$. Then clearly $n=1=r$ and thus $n-m+r = 1-0+1 = 2$, as desired.

$m-1 \rightarrow m$: Assume we established the claim for all graphs

with less than m many edges. Consider G with m many edges.

If G is a tree, then we know that $n=m+1$ and $r=1$ (whence $n-m+r = m+1-m+1 = 2$, as desired).

Otherwise, G contains at least one cycle. Let $e \in E(G)$ be one edge on that cycle. Then by Observation 5.10, e is a bound for two regions R_i and R_j . Note that in $G-e$ the regions R_i and R_j merge together to one new region R'_i , whence $G-e$ has one less region than G . Thus, by I.H. we get

$$\|G-e\| - \|G-e\| + (r-1) = 2 = n - (m-1) + (r-1) = n-m+r, \text{ as desired. } \square$$

5.13 Remark

1) Every subgraph of a planar graph is planar.


2) If G is planar and R is any region, then in the induced planar representation for $B(R)$ (i.e. deleting all drawings apart from $B(R)$) R is still a region and $B(R)$ its boundary.

\Rightarrow 3) For any region we have $b(R) \in \{0\} \cup \mathbb{N}_{\geq 3}$. In particular, if

$$b(R)=3, \text{ then } B(R) \cong C_3.$$

5.2 Nonplanar Graphs

Imagine you want to connect three houses each to gas, electricity and water, s.t. the pipes don't intersect. This question boils down to asking: Is



(This is why $K_{3,3}$ is also called the utility graph.)

5.14 Theorem

The utility graph $K_{3,3}$ is not planar.

Proof

Starting for a contradiction, assume $K_{3,3}$ was planar. Then it needed to have $r = ||K_{3,3}|| - |K_{3,3}| + 2 = 9 - 6 + 2 = 5$ regions. On the other hand, as $K_{3,3}$ is bipartite and thus does not contain odd cycles, every region has at least four bounds by

Remark 3.13, 3). Hence $\sum_{i=1}^5 b(R_i) \geq 5 \cdot 4 = 20 > 2 \cdot 9 = 2 \cdot ||K_{3,3}||$, contradicting Fact 5.9.

Thus, $K_{3,3}$ cannot be planar. \square

5.15 Theorem

Let G be planar of order at least 3. Then $|E| \leq 3(|V| - 2)$.

Further, if equality holds then every region is bounded by exactly three edges.

Proof

Assume G consists of k connected components. The equation clearly holds for forests as then $|E| = |V| - k \leq 3|V| - 6 \iff 6 - k \leq 6 \leq 2|V|$.

Otherwise, $r \geq 2$ and G contains a cycle. Recall that

$$(*) 3 \cdot r \leq \sum_{i=1}^r b(R_i) \leq 2 \|G\|. \text{ Hence, by Euler's generalised formula,}$$

we get $r = \|G\| - |G| + k + 1 \geq \|G\| - |G| + 2$. This yields

$$2 \|G\| \geq 3r \geq 3(\|G\| - |G| + 2), \text{ whence } 3(|G| - 2) \geq \|G\|, \text{ as desired.}$$

Finally, for equality to hold we need in particular that


$$2 \|G\| = 3r, \text{ whence also } 3r = \sum_{i=1}^r b(R_i). \text{ Thus, every region}$$

has boundary degree 3 as desired. \square

5.16 Corollary

The complete graph K_n is planar iff $n \leq 4$.

Proof

First note that $K_4 =$  is planar, and hence so are its subgraphs K_1, K_2 and K_3 .

Now, for K_5 , by Lemma 5.15 we get $10 = |K_5| \leq 3(\|K_5\| - 2) = 3(3) = 9$, a contradiction. Hence, K_5 is not planar and so is neither of the K_n for $n \geq 5$, as they contain K_5 as a subgraph. \square

5.17 Theorem

If G is planar, then $\delta(G) \leq 5$.

Proof

Let G be a planar graph and set $n = |G|$ and $m = \|G\|$. Aiming for a contradiction, assume $\delta(G) > 5$. Then in particular, $n \geq 6$.

$$\text{Then } 6 \cdot n \leq \sum_{v \in V_G} \deg(v) \stackrel{(1.13)}{=} 2m \stackrel{(5.15)}{\leq} 2(3 \cdot (n-2)) = 6n - 12 \text{ yields}$$


a contradiction. \square

5.3 Kuratowski's Theorem

We have seen that the graphs $K_{3,3}$ and K_5 are not planar. It turns out that these two graphs are the major obstruction for any graph to be planar. This section gives an introduction to Kuratowski's theorem. But first, we want to introduce a new notation.

5.18 Example

Consider



Note that neither of G or H contains K_5 (or $K_{3,3}$) as a subgraph. Nevertheless, they both look so similar to K_5 that they should not be planar.

Indeed, G arises from K_5 by replacing the edge xz by $\{xy, yz\}$ and H arises from G by replacing the edges $\{xz, ad\}$ by $\{xy, yz, ab, bc, cd\}$. This process is called subdivision.


5.19 Definition

- 1) Let G be a graph and $e = xy \in E(G)$. An **edge subdivision** of e is the replacement of e by a finite path of length ≥ 2 starting in x and ending in y .
- 2) Let G and H be graphs. Then H is called a **subdivision** of G iff H can be obtained through a finite sequence of edge subdivisions.

5.20 Examples

1) The graphs G, H from 5.18 are edge subdivisions of K_5 .

2) The graph



is not an edge subdivision of K_5 , as the edges $\{xy, yz\}$ were added rather than replacing the edge xz .

5.21 Lemma

A graph G is planar iff every subdivision of G is planar.

Proof

" \Rightarrow " Consider a planar representation Γ of G . Let H be a subdivision of G where a sequence of n -many edge subdivisions were performed. We do induction on n .

If $n=0$, then $H=G$ is clearly planar.

Now assume we proved the claim for n and consider H arising from G through $n+1$ many subdivisions. Let H_0 be the graph arising from G through the first n -many subdivisions. By 1.b. H_0 is planar and H can be obtained from H_0 through exactly one edge-subdivision. Let Γ_0 be a planar drawing of H_0 and $e=xy$ be the edge which is subdivided to obtain H , say by replacing it by the path $p=(x=x_0, x_1, \dots, x_k=y)$ for $k \geq 2$. Let \vec{xy} be the geodesic from x to y in \mathbb{R}^2 . Then we draw in the vertex x_i at the point $x + \frac{i}{k}\vec{xy}$ for any i . This yields a planar representation of H , as desired. □

5.22 Corollary

If G contains a subdivision of $K_{3,3}$ or K_5 as a subgraph, then G is not planar. (Or: If G planar \Rightarrow no subdivision of $K_5, K_{3,3} \subseteq G$)

5.23 Theorem (Kuratowski's Theorem)


A graph is planar if and only if it does not contain a subdivision of $K_{3,3}$ or K_5 as a subgraph.

5.24 Remark


The left over, hard direction is the backwards direction, i.e. if G does not contain a subdivision of $K_{3,3}$ or K_5 , then G is planar.

INTERMEZZO

Prove that the Peterson graph is not planar.



Note that the subgraph $G - \{x_2, x_{10}, x_3, x_4\}$ is a subdivision of $K_{3,3}$ given through



and the edge subdivisions

$$x_5 x_9 \rightarrow \{x_5 x_4, x_4 x_9\}$$

$$x_2 x_8 \rightarrow \{x_2 x_3, x_3 x_8\}$$

$$x_5 x_8 \rightarrow \{x_5 x_{10}, x_{10} x_8\}$$

$$x_2 x_9 \rightarrow \{x_2 x_7, x_7 x_9\}$$

and

Chapter 6 - Graph Colourings

Disclaimer

The title is actually somewhat misleading, as our first convention will be that we can understand different colours simply as different positive integers. Hence we will use the "colours" $1, 2, 3, \dots$ instead of red, blue, brown etc. Visualisations can nevertheless reassociate these numbers with colours.

6.1. The chromatic number

6.1 Definition

Let G be a graph. A k -colouring is a function $K: V_G \rightarrow \{1, 2, \dots, k\}$ for $k \in \mathbb{Z}_+$ s.t. $K(x) = K(y)$ implies $xy \notin E_G$, i.e. adjacent vertices receive distinct colours. We call $\{1, 2, \dots, k\}$ the colours of K . We say that K is a colouring of G if it is a k -colouring for some k and we call G k -colourable if there is a k -colouring for G .

6.2 Remarks

Let G be any graph.

- 1) If G is k -colourable, then it is j -colourable for any $j \geq k$.
- 2) If K is a k -colouring of G , then $V_r := K^{-1}(r) = \{v \in V_G \mid K(v) = r\}$ is an independent subset of G for any $r \in \{1, 2, \dots, k\}$.

3) Every graph of order n is clearly n -colourable as for

$V_G = \{v_1, v_2, \dots, v_n\}$ we can simply define the injective function

$k: V_G \rightarrow \{1, 2, \dots, n\}$ via $k(v_i) = i$. Any injective such function is clearly a colouring.

\Rightarrow 4) Remarks 2) and 3) indicate that the interesting question will be to find small k s.t. G is k -colourable

6.3 Example

- Consider the graph C_5 . Let's try to find a minimal k s.t. G is k -colourable.


\Rightarrow It seems we need at least

3-colours. C_5 is 3-colourable, as $k: V_G \rightarrow \{1, 2, 3\}$

via $k(v_1) = k(v_4) = 1$, $k(v_2) = k(v_5) = 2$ and $k(v_3) = 3$ is a 3-colouring

for G . We also could have defined k via $V_1 = \{v_1, v_4\}$, $V_2 = \{v_2, v_5\}$

and $V_3 = \{v_3\}$.



6.4 Remarks

- Note that every k -colouring k of G gives rise to a partition of V_G into sets V_1, V_2, \dots, V_k s.t. no two vertices in V_i are adjacent. G is hence k -partite. Note that V_i could be empty for some i .

- As each V_i is an independent set, we get $k \geq \frac{|G|}{\alpha(G)}$ (H.W.).

6.5 Definition

The chromatic number $\chi(G)$ of a graph G is the smallest positive integer $k \in \mathbb{Z}_+$ s.t. G is k -colourable.

6.6 Examples

- 1) $\chi(G)=1$ iff $G = E_n$ is the empty graph for some n .
- 2) $\chi(K_n) = n$, as all of the n -many vertices are pairwise adjacent.
- 3) $\chi(K_{n,m}) = 2$, as we can colour each part of the partition in one colour.
- 4) $\chi(P_n) = 2$ for $n \geq 2$.
- 5) $\chi(C_n) = \begin{cases} 2 & \text{if } n \text{ is even} \\ 3 & \text{otherwise} \end{cases}$ (Hw)

6.7 Remark

- 1) G is k -colourable iff $k \geq \chi(G)$.
- 2) $\frac{|G|}{\alpha(G)} \leq \chi(G) \leq |G|$ for any graph G .
- 3) G is 2-colourable iff G is bipartite (iff $\chi(G) \leq 2$)
- \Rightarrow 4) If F is a forest, then $\chi(F) = 2$.
- 5) If $H \subseteq G$ is a subgraph of G , then $\chi(H) \leq \chi(G)$.

6.8 Definition


Let $n_1, \dots, n_k \in \mathbb{Z}_+$ be positive integers, $k \geq 1$. The complete k -partite graph K_{n_1, n_2, \dots, n_k} is the graph whose vertex set is the disjoint union of k -many pairwise disjoint sets V_1, \dots, V_k with $|V_i| = n_i$; and edge set $E = \{uv \mid u \in V_i, v \in V_j, i \neq j\}$. If we don't care about the specific value of k , we call K_{n_1, \dots, n_k} the complete multipartite graph.

6.9 Example

The complete 3-partite graph $K_{3,1,2}$ is

We see that even for small numbers, the graph is

difficult to draw. For which n_1, n_2 is K_{n_1, n_2} planar?



6.10 Lemma

Let G be a graph and $k \geq 1$. Then G is k -colourable iff G is a subgraph of a complete k -partite graph.

Proof

G is k -colourable

iff ex. $K: V_G \rightarrow \{1, 2, \dots, k\}$ s.t. $K(v) = k(v) \rightarrow uv \notin E_G$

iff ex. $K: V_G \rightarrow \{1, 2, \dots, k\}$ s.t. $K^{-1}(i)$ is an independent set (or empty) for all i

iff we can partition V_G into at k -many independent sets V_1, \dots, V_k ($V_i = \emptyset$ allowed)

iff G is a subgraph of K_{n_1, \dots, n_k} for some $n_1, \dots, n_k \in \mathbb{Z}_+$.



Next we want to introduce a greedy algorithm for finding a vertex colouring of a given graph. This algorithm is more efficient than giving every vertex a different colour, but it does not always produce a $\chi(G)$ -colouring.

6.11 The Greedy Algorithm

Let G be a graph of order n .

1) Label the vertices by v_1, \dots, v_n .

2) Fix the set of available colours to be $\{1, 2, \dots, n\}$.

3) Let $i = 1$.

4) While $i \leq n$




- Colour v_i with the smallest available colour not used on any of its previously coloured neighbours, i.e. set

$$k(v_i) := \min \{1, 2, \dots, n\} \setminus \{k(v_j) \mid j < i, v_j \in N(v_i)\}.$$

- set $i \rightarrow i+1$.

INTERESTING

Let  = 1,  = 2,  = 3,  = 4



Greedy algorithm yields
a 2-colouring.

Greedy algorithm yields
a 3-colouring.

Greedy algorithm yields
a 4-colouring.

\Rightarrow The colours used depend heavily on the labeling.

6.12 Lemma

Let G be a graph and K be a colouring function produced by the greedy algorithm applied to G . Then for all $v \in V_G$, we get

$$K(v) \leq \deg(v) + 1.$$

Proof

We do induction on $|G|=n$. For $n=1$, clearly $G=K_1$,

and $\chi(v_1) = 1 = \deg(v_1) + 1$. Now assume the claim holds

for any graph of order at most n and consider G with $|G|=n+1$.

Run the greedy algorithm on G . After the first n rounds, we

obtain a "greedy" colouring of $G - v_{n+1}$, whence

$$\chi(v_i) \leq \deg(v_i) + 1 \leq \deg(v_i) + 1 \text{ for all } 1 \leq i \leq n. \text{ Now we run the}$$

last round and assign $\chi(v_{n+1}) = \min \{1, \dots, n\} \setminus \{\chi(v_j) \mid v_j \in N(v_{n+1})\}$.

At most $\deg(v_{n+1})$ -many colours have been used on its neighbours, whence among $\{1, 2, \dots, \deg(v_{n+1}), \deg(v_{n+1}) + 1\}$, there is at least one colour available. As the algorithm picks the smallest available colour, we conclude $\chi(v_{n+1}) \leq \deg(v_{n+1}) + 1$, as desired. □

6.13 Corollary

For any graph G we have $\chi(G) \leq \Delta(G) + 1$.

Proof

Apply the greedy algorithm on G . Then for any $v \in V_G$ we get

$$\chi(v) \leq \deg(v) + 1 \leq \Delta(G) + 1, \text{ whence } \chi: V_G \rightarrow \{1, 2, \dots, \Delta(G) + 1\}$$

$$\text{and } \chi(G) \leq \Delta(G) + 1. \quad \square$$

Finding the chromatic number of a graph is a very important, but computationally hard problem. It is known to be NP-hard and its best runtime is in $\Theta(2^n)$. We hence need to establish good bounds to approach the problem effectively.

6.14 Remark

- For any graph G we have $\chi(G) \leq \Delta(G) + 1$.
- The bound is sharp, as
 - $\chi(K_n) = n = (n-1) + 1 = \Delta(K_n) + 1$ for all $n \in \mathbb{Z}_+$ and
 - $\chi(C_n) = 3 = 2 + 1 - \Delta(C_n) + 1$ for all odd $n \in \mathbb{Z}_+$.
- But are there more examples to witness that the bound is sharp (i.e. cannot be improved)? No.
- Note: What do C_n and K_n have in common? They are regular!

Goal of this lecture:

6.15 Theorem (Brooks's Theorem, 1941)

If G is a connected graph which is neither complete, nor an odd cycle, then $\chi(G) \leq \Delta(G)$.

We will split the proof into several Lemmata. Set $\Delta(G) := 1$.

6.16 Step 1 - Remark on $\Delta \leq 2$

Theorem 6.15 holds for $\Delta \leq 2$, as if $\Delta=0$, then $G=k$, and if $\Delta=1$, then $G=k_2$ is complete, which is excluded.

For $\Delta=2$ either $\delta(G)=\Delta=2$ and G is an even cycle, whence $\chi(G)=2 \leq \Delta(G)$ holds, or $\delta(G)=1$ and G is a path of length at least 2, and again $\chi(G)=2 \leq \Delta(G)$.

Thus, from now on, consider such G which satisfy $\Delta(G) \geq 3$.

6.17 Step 2 - Lemma on non-regular graphs.


If G from 6.15 is not regular, then $\chi(G) \leq \Delta(G)$ (i.e. the theorem holds).

Proof

If G is non-regular, then ex. $v \in V_G$ s.t. $\deg(v) < \Delta(G)$.

The idea is to introduce a smart colouring which uses at most $\Delta(G)$ many colours. Note that as G is (finite and) connected, $\text{ecc}(v) = k < \infty$. Let $S_i := \{u \in V_G \mid d(u, v) = i\}$. Hence, $S_0 = \{v\}$ and $S_k = \{u \in V_G \mid d(u, v) = \text{ecc}(v)\}$. Note further that for $1 \leq i \leq k$ every vertex $u \in S_i$ has at least one neighbour in S_{i-1} (i.e. the predecessor of u on a vu -path of shortest length).

We get:



We want to apply the greedy algorithm after labelling the vertices of G in a smart way: We start by randomly labelling the vertices in S_0 and then those in S_{k-1} and so on and so forth until the last vertex v gets labelled v_n where $n = |G|$.

Hence, if $v_i \in S_\ell$ and $v_j \in S_m$ for $\ell < m$, then $i > j$.

Now we run the greedy algorithm and observe which colours the vertices get.

For $v_i \in S_\ell$ with $\ell \neq 0$ (i.e. $v_i \neq v_n = v$), note that v_i gets the smallest available colour not used on its previously coloured neighbours, i.e. on its neighbours in $S_\ell \cup S_{\ell+1} \cup \dots \cup S_k$.

As v_i has at least one neighbour in $S_{\ell-1}$, these are at most $\deg(v_i) - 1 \leq \Delta(G) - 1$ many. Hence, at least

one of the colours $\{1, 2, \dots, \Delta(G)\}$ is still available whence $k(v_i) \leq \Delta(G)$ as desired.

For $v_n = v$, it gets the smallest available colour not used on any of its neighbours, as it is the last to get coloured. But as by choice, $\deg(v_n) < \Delta(G)$, again one of the colours $\{1, 2, \dots, \Delta(G)\}$ must be available and $k(v_n) \leq \Delta(G)$.

Hence, $k(u) \leq \Delta$ for all $u \in V_0$, whence G is Δ -colourable and $\chi(G) \leq \Delta$, as desired. \square

INTERMEzzo

Consider the set V of all courses offered at AUC.

We want to schedule finals s.t. (finally!!) no student has to take two exams on the same day.


Can you find a graph structure on V and use graph theoretic tools to determine how many exam dates we need and which courses can have simultaneous exams?

6.18 Step 3 - Regular graphs with cut vertices

Let G be a connected, regular graph with $\Delta(G) \geq 3$. If G contains a cut vertex (i.e. $k(G) = 1$), then $\chi(G) \leq \Delta(G)$.

6.19 Example

You might wonder if these graphs exist. Here is an example:



Proof of 6.18

Let v be a cut vertex in G , i.e. $G-v$ is disconnected. Let H_1, H_2, \dots, H_k be the distinct connected components of $G-v$ with $k \geq 2$. Further, define $G_i := \langle V(H_i) \cup \{v\} \rangle$ to be the graph induced on H_i with v . Note that $\deg^{G_i}(v) < \deg^G(v) = \Delta$, while still $\deg(u) = 1$ for all other $u \in V(G_i)$. Hence $\Delta(G_i) = \Delta$. By Step 2, we can find Δ -colouring of G_i . Possibly after permuting the colours, we may assume that $k_i(v) = k_j(v)$ for all $1 \leq i, j \leq k$. Hence $K = k_1 \cup \dots \cup k_k$ is the desired Δ -colouring of G . □

6.20 Auxiliary Step 4 - Fact

Let G be a regular, non-complete, 2-connected graph with $\Delta(G) \geq 3$. Then there exist vertices v_1, v_1, v_2 s.t. $v_1, v_2 \in N(v_1)$, but v_1 is not adjacent to v_2 and additionally $G - \{v_1, v_2\}$ is still connected.

6.21 Step 5 - Left overs


Let G be regular, 2-connected, non-complete with $\Delta(G) \geq 3$.

Then $\chi(G) \leq \Delta(G)$.

Proof

By Fact 6.20, there exist vertices v_1, v_1, v_2 s.t. $v_1, v_2 \in N(v_1)$, $v_1 \notin N(v_2)$ and $G - \{v_1, v_2\}$ is connected. Let $H := G - \{v_1, v_2\}$ and as in step 1, partition V_H via $S_i = \{u \in H \mid d(v_1, u) = i\}$. Let $|S_i| = n_i$ and label all vertices as in step 1, except that the labels of v_1, v_2 and v_1 stay. So if $\text{ecc}^H(v_1) = k$, we start labeling the vertices in S_k with v_3, v_4, \dots and so on, followed by the vertices in S_{k-1}, \dots until we reach

$S_0 = \{v_n\}$, which is already labeled. Now we get the picture



We are ready to run the greedy algorithm.

First, we get $\chi(v_1) = 1$ and then also $\chi(v_2) = 1$, as they are not adjacent.

Now, any v_i with $3 \leq i < n$, there is at least one neighbour which is not coloured in step i . Precisely, if $v_i \in S_{i-1}$, then there is a neighbour $v_j \in S_{i-1}$ with $i < j$ which is not coloured. Its $\deg(v_i) = \Delta$, at least one of the colours $\{1, 2, \dots, \Delta\}$ is still available, whence $\chi(v_i) \leq \Delta$.

Finally, $\deg(v_n) = \Delta$ and all its neighbours have been coloured, but two of its neighbours, v_1 and v_2 , have the same colour, whence once again, at least one of the colours $\{1, 2, \dots, \Delta\}$ is still available and also $\chi(v_n) \leq \Delta$.

We hence proved that $\chi(G) \leq \Delta$, as desired. □

This concludes the proof of Brooks's Theorem. The only blackbox we are using is Fact 6.20, due to time reasons. You can find the proof in the lecture notes of Dr. Daoud.

6.22 Corollary

Let G be a graph with connected components H_1, H_2, \dots, H_k ,

1) Then $\Delta(G) = \max \{\Delta(H_i) | 1 \leq i \leq k\}$ and $\chi(G) = \max \{\chi(H_i) | 1 \leq i \leq k\}$.

$\Leftrightarrow 2) \chi(G) = \Delta(G) + 1$ if and only if ex. i s.t. H_i is either a complete graph or an odd cycle and $\Delta(H_i) = \Delta(G)$.

6.23 Homework

Let G be a graph of order n , then


$$\frac{n}{\alpha(G)} \leq \chi(G) \leq n + 1 - \alpha(G).$$

The rest of the lecture is devoted to give a last bound for the chromatic number of a graph.

6.24 Definition

The clique number of a graph G , denoted by $\omega(G)$, is the largest positive integer m s.t. G contains K_m as a subgraph.

6.25 Example



Then $\omega(G) = 3$ witnessed
by $\langle \{v_1, v_2, v_3\} \rangle \cong K_3$

$\omega(H) = 4$ witnessed
and
by $\langle \{v_1, v_2, v_3, v_4\} \rangle \cong K_4$
and H being planar.

6.26 Lemma


For any graph G we get $\omega(G) \leq \chi(G)$.

Let $\omega(G) = l$, then G contains K_l as a subgraph which implies that we already need l colours just to colour K_l and $l = \omega(G) \leq \chi(G)$.

The question arises whether $\omega(G)$ -many colours should always be enough to colour a graph G . This hope rather quickly fails, as for example $\omega(C_5)=2$, but $\chi(C_5)=3 > \omega(G)$. Another example is given below.

6.27 Example

Let G be the graph consisting of the disjoint union of C_5 and C_3 together with all edges between C_3 and C_5 , i.e.



Then $\omega(G)=5$, as e.g.

$\langle \{v_1, v_2, v_3, v_4, v_5\} \rangle \cong K_5$ (so $\omega(G) \geq 5$),

but any 6 vertices need to include

at least three vertices from C_5 , which cannot

be mutually incident, so $\omega(G) < 6$, and thus $\omega(G)=5$.

Further, $\chi(G)=6$: The cycle C_5 and C_3 each need at least 3 colours, and these colours have to be distinct as the vertices are mutually incident.

Let us summarize all bounds that we established:

Summary

Let G be any graph of order n .

1) $\chi(G) \leq \Delta(G) + 1 \leq n$.

2) If G is connected then $\chi(G) \leq \Delta(G)$ iff G is neither complete nor an odd cycle.

3) $\omega(G) \leq \chi(G)$ and $\frac{n}{\Delta(G)} \leq \chi(G) \leq n + 1 - \alpha(G)$.

6.2 - The 4-Colour Problem


6.28 - The problem (Francis Guthrie, 1852)

Given any map in the plane - how many colours do we need to colour it in a way s.t. no countries who share part of their border (more than a point) have the same colour?


We can rephrase this in graph theoretic terms. Note that we can represent the issue as a graph where the vertices represent countries and edges connect countries with touching borders. Such a graph will be planar.

6.29 Example


The map




can be represented by the planar graph:



Now, a colouring of the map exactly corresponds to a vertex-colouring of the associated graph, e.g.



yields



For this graph, we obtained a 4-colouring. We know that

$\omega(G) \leq \chi(G)$ for any graph and $\omega(G) \leq 4$ for any planar graph.

But are 4 colours always enough?

6.30 The Four-Colour-Theorem (1976, Appel, Haken)

If G is planar, then $\chi(G) \leq 4$. i.e. every planar graph is 4-colourable.

6.31 Some history

- 1852 - Problem introduced by Francis Guthrie to DeMorgan (his prof)
 - 1852-1879 Brilliant minds, incl. Hamilton, Cayley, Peirce... tried to solve the problem without success
 - 1879 - Alfred Kempe announced a proof
 - 1890 - Fatal mistake was found in proof
- soon after - Heawood + Kempe prove that $\chi(G) \leq 5$.
- 1976 - After 124 years a proof was presented by Appel-Haken
 - ↳ relies heavily on computers, not accepted by all mathematicians
 - 1996 - Easier proof presented, but still uses computers.

→ The search continues.

Maybe somewhat surprisingly, the proof for $\chi(G) \leq 5$ is much more accessible. Recall that if G is planar, we have $\delta(G) \leq 5$. (Thm. 5.17)

6.32 Theorem

If G is planar, then $\chi(G) \leq 5$, i.e. G is 5-colourable.

Proof


Let $|G|=n$. We proceed by induction on n .

If $n \leq 5$, the theorem clearly holds as $\chi(G) \leq |G|$ for any G .

$n \rightarrow n+1$: Assume any planar graph of order at most n

is 5-colourable and consider G s.t. $|G|=n+1$. As $\delta(G) \leq 5$, there exists $v \in V_G$ s.t. $\deg(v) \leq 5$. Consider the planar graph $G-v$. By L.H., $G-v$ is 5-colourable. Let k be that 5-colouring. Now, if among the neighbours of v at least one of $\{1, 2, 3, 4, 5\}$ was not used, we can use that colour for v and obtain the desired 5-colouring of G .

Otherwise, $\deg(v) = 5$ and all neighbours $\{w_1, w_2, w_3, w_4, w_5\}$ of v are coloured in a different colour, e.g. $k(w_i) = i$. We obtain:




as a subgraph of G .

Now, we make a case

distinction.

Case 1: Assume there is no w, w_3 -path that entirely uses vertices coloured with colours 1 and 3. Then let H be the subgraph of G containing all paths that start in w_1 and use only vertices of colour 1 and 3, i.e. $H = \cup \{p \mid p \text{ is a } w, u\text{-path} \& k(x) \in \{1, 3\} \forall x \in p\}$.

Note that $w_1 \notin H$, but $w_3 \notin H$ by assumption.




Now, in H , exchange the colours 1 and 3 and observe that that new colouring \tilde{k} is still a valid colouring for G , but now $\tilde{k}(w_1)=3$. To see that it is valid, assume for contradiction that $\exists x, y \in E_G \quad \tilde{k}(x)=\tilde{k}(y)$ and $x, y \in H$. Then wlog we may assume $x \in H$, $y \notin H$ and

$\tilde{k}(x) = \tilde{k}(y) = 1$. But then $k(x) = 3$, $k(y) = 1$ and there is a $\{1,3\}$ -coloured w, x -path p . Finally, $p^*(y)$ would be a $\{1,3\}$ -coloured w, y -path, whence $y \in H$, contradicting our assumptions.


Hence, \tilde{k} is a 5-colouring of $G-v$, but now the neighbours of v only use colours $\{2,3,4,5\}$, whence we can set $\tilde{k}(v) = 1$ and obtain the desired 5-colouring of G .

Case 2: Assume there is a $\{1,3\}$ -coloured $w_1 w_3$ -path p in $G-v$.

as G is planar, we have two options:



or:



In either case, any $w_2 w_4$ -path now would have to contain at least one vertex from p , as G is planar. But this means that there is no $w_2 w_4$ -path which only uses colours 2 and 4.

We can hence apply Case 1 to w_2 and w_4 and obtain the desired 5-colouring of G . □

INTERMEZZO

Play the C_3 -Game! One player is red, one is blue

In alternating turns, colour the edges of 1) K_5 and then 2) K_6 .

in your colour. You win if you are the first to produce a

monochromatic triangle! What is the difference of playing it on K_5 vs. K_6 ?

6.3 Chromatic Polynomials

History


- Introduced 1912 by Georg David Birkhoff to tackle the 4-colour problem (he hoped for a negative answer)
- $P_G(k)$ shall describe the number of k -colourings of G
- These will be polynomials in k of degree $|G|=n$, i.e. $P_G(k) = a_n k^n + \dots + a_1 k + a_0$.
- Birkhoff hoped to use strong tools from analysis and algebra to find roots of these polynomials
- In particular, he hoped to find a planar graph G , which has $k=4$ as a root, i.e. $P_G(4)=0$, whence G has no 4-colouring, whence the 4-colour theorem would be wrong
- Even though we know that he had no chance of success, he still developed many tools which are crucial in the area of algebraic graph theory:
⇒ Sometimes truly the way is the goal.

6.3.3 Definition


- 1) Let k_1 and k_2 be two colourings of the same graph G . We say that k_1 is different from k_2 ($k_1 \neq k_2$) if there is some $v \in V_G$ s.t. $k_1(v) \neq k_2(v)$.
- 2) We denote by $P_G(k)$ the number of different k -colourings of G .

6.34 Example

Consider the following colourings of K_4

and  . (k') Then K and K' are

different, as $k(v_3) = 3 \neq 4 = k'(v_3)$.



(k)

6.35 Discussion

- How many k -colourings of K_4 are there?

↳ We can choose any of the 4 colours to colour v_1 , then any of the remaining 3 for v_2 and so on. In the end we obtain $4 \cdot 3 \cdot 2 \cdot 1 = 4!$ many 4-colourings.

- What about 6-colourings?

↳ Following the same thoughts, we obtain $6 \cdot 5 \cdot 4 \cdot 3 = \frac{6!}{(6-4)!}$ many 6-colourings.

- And 3-colourings?

↳ As $\chi(K_4) = 4$, there are no 3-colourings.

6.36 Definition

Let G be any graph. Then we denote by $P_G(k)$ the number of possible different colourings using at most the colours $\{1, 2, \dots, k\}$.

6.37 Remark

- Generalising our discussion above, we obtain

$$P_{K_n}(k) = \begin{cases} 0 & \text{if } k < n \\ \frac{k!}{(k-n)!} & \text{if } n \leq k \end{cases}$$

- Further check quickly that $P_{E_n}(k) = k^n$.

6.38 Remark

The following equivalent for any graph G and $k \in \mathbb{Z}_+$.

- 1) $P_G(k) \geq 1$
- 2) $\chi(G) \leq k$
- 3) G is k -colourable.

In order to show that $P_G(k)$ is a polynomial of degree k , we need an important observation which will allow us to use inductive arguments. To make sense of it, we need the following definition.


6.39 Definition (Edge Contraction)

Let G be a graph and $e \in E(G)$, say $e=uv$. Then the **edge contraction** G/e is the graph obtained from G by the following:

- 1) Delete the edge e from G , i.e. construct $G-e$.
- 2) Identify the vertices u and v as one new vertex denoted uv .
- 3) leaving only one copy of any resulting multi-edges.

6.40 Example

Consider G and $e \in E(G)$ as follows:



6.41 Theorem (Birkhoff-Lewis, 1946)

Let G be any non-empty graph and $e=uv \in E(G)$. Then

$$P_G(k) = P_{G-e}(k) - P_{G/e}(k).$$

Proof

Claim 1: $P_{G/e}(k)$ is equal to the number of k -colourings of $G-e$, which assign the same colour to u and v .

↪ Let K be a k -colouring of $P_{G/e}$. Then we can create a k -colouring \tilde{K} of $G-e$ via $\tilde{K}(x) = \begin{cases} K(uv) & \text{if } x=u \text{ or } x=v \\ K(x) & \text{otherwise} \end{cases}$. Hence, there are at least as many k -colourings of $G-e$ which assign the same colour to u and v as there are k -colourings of G/e . On the other hand, if \tilde{K} is any k -colouring of $G-e$ which assigns the same colour to u and v , then we can define a new k -colouring K of G/e by setting $K(x) = \begin{cases} \tilde{K}(u) & \text{if } x=uv \\ \tilde{K}(x) & \text{else} \end{cases}$. Hence, there are at least as many k -colourings of G/e as there are k -colourings of $G-e$ assigning the same colour to u and v .

Claim 2: There are as many k -colourings of G as there are k -colourings of $G-e$ assigning different colours to u and v .

↪ If \tilde{K} is a colouring of G , then clearly it is a k -colouring of $G-e$ assigning different colours to u and v and vice versa. (Note that both graphs have the same vertex set).

$$\begin{aligned} \text{Now, } P_{G-e}(k) &= |\{K \text{ } k\text{-colouring of } G-e\}| \\ &= |\{K \text{ } k\text{-colouring of } G-e \text{ with } K(u)=K(v)\}| + \\ &\quad + |\{K \text{ } k\text{-colouring of } G-e \text{ with } K(u) \neq K(v)\}| \\ &= P_{G/e}(k) + P_G(k). \end{aligned}$$

Thus, $P_G(k) = P_{G-e}(k) - P_{G/e}(k)$, as desired.


INTERMEZZO

compute $P_{C_3}(k)$ for any k - first by combinatorial arguments

- then by using Birkhoff Lewis

(Use that $P_{E_n}(k) = k^n$, reduce to that)


Solution




1) We have k -many choices to colour u , then all $(k-1)$ many left over colours for v and finally $(k-2)$ many choices for w , so $k \cdot (k-1)(k-2) = k^3 - 3k^2 + 2k$.
Here, this does not depend on the order of choosing u, v, w .

2) Birkhoff-Lewis:


• Let $G_0 = G - e_1$, $G_1 = G/e_1$, i.e. G_0



• Now let $G_{00} := G_0 - e_2$, $G_{01} := G_0/e_2$, $G_{10} := G_1 - e_2 \wedge e_3$, $G_{11} := G_1/e_2 \wedge e_3$,



Finally, let $G_{000} := G_{00} - e_3$, $G_{001} := G_{00}/e_3$, i.e. G_{000}



as well as $G_{010} := G_{01} - e_3$ and $G_{011} := G_{01}/e_3$, i.e. G_{010}




We know the chromatic polynomial for the empty graph $P_{E_n}(k) = k^n$. Thus

$$\begin{aligned}
 P_{C_3}(k) &= P_{G_0}(k) - P_{G_1}(k) \\
 &= P_{G_{00}}(k) - P_{G_{01}}(k) - P_{G_{10}}(k) + P_{G_{11}}(k) \\
 &= P_{G_{000}}(k) - P_{G_{001}}(k) - P_{G_{010}}(k) + P_{G_{011}}(k) - P_{G_{10}}(k) + P_{G_{11}}(k) \\
 &= k^3 - k^2 - k^2 + k - k^2 + k = k^3 - 3k^2 + 2k.
 \end{aligned}$$

6.42 Remark

• Combinatorics seems faster than Birkhoff-Lewis, so why bother?

- 1) We do not need to always reduce back to E_n , once we know the chromatic polynomial of other graphs \rightarrow it becomes much faster.
- 2) It allows inductive arguments in proofs (see below).
- 3) Combinatorial thoughts depend on the choice of vertices and hence do not always give the correct number!



e.g.: $G = \dots P_G(3) \dots$ there are 3 colours to pick from for v_5 , also 3 for v_4 , 2 for v_1 , 2 for v_3 and either 1 or 0 for $v_2 \dots$ we need a case distinction!
 \rightsquigarrow Things get messy very quickly.

But: Applying Birkhoff-Lewis, we always get a solid solution!

Note: $P_{C_3}(3) = 3^3 - 3 \cdot 3^2 + 2 \cdot 3 = 6$. Now set $e_1 := v_1v_4$, $e_2 := v_2v_5$.

Then $G_0 := G - e_1$ is

, $G_1 := G/e_1$ is

Further $G_{00} := G_0 - e_2$ is

, $G_{01} := G_0/e_2$ is

and $G_{10} := G_1 - e_2$ is

, $G_{11} := G_1/e_2$ is

$$\begin{aligned} \text{Now } P_G(3) &= P_{G_0}(3) - P_{G_1}(3) = P_{G_{00}}(3) - P_{G_{01}}(3) - P_{G_{10}}(3) + P_{G_{11}}(3) \\ &= 3 \cdot 3 \cdot 6 - 3 \cdot 6 - 3 \cdot 6 + 6 \\ &= 6(9 - 3 - 3 + 1) = 24. \end{aligned}$$

6.43 Theorem

Let G be a graph of order n . Then the following hold.

- 1) $P_G(k)$ is a polynomial in k of degree n , $P_G(k) = a_n k^n + \dots + a_1 k + a_0$.
- 2) We always have $a_n = 1$ and $a_0 = 0$, i.e. $P_G(k) = k^n + a_{n-1} k^{n-1} + \dots + a_1 k$.
- 3) The a_i alternate in sign and $a_{n-i} = -|E(G)|$.


Chapter 7 - Matchings

7.1 Motivation


Imagine a group of people has to split up in pairs and take part in a competition. Every person has several people they'd be happy to compete with, and we assume this is symmetric.

This situation can be modeled using a graph. How would a possible matching look like which would please everyone?

Example:



Possible matchings are



If we pair
2 with 5
3 and 4 with 1,
this is still a matching.

But we have 2 happy teams and
can't form another one. It is not perfect...

Observation: The goal will be to pick a set of edges which do not share end vertices.

7.2 Definition

Let G be any graph.

- i) A matching for G is a set $M \subseteq E(G)$ of pairwise disjoint edges.
- ii) $v \in V(G)$ is called M -saturated if exists $e \in M$ st. $v \in e$ (i.e. if it is the endpoint of some edge in M). Otherwise,

we call ν M-unsaturated.


iii) We call a matching M maximal iff $M \cup \{e\}$ is not a matching for any $e \in E(G) \setminus M$.

iv) We say that M is a maximum matching if it has the largest cardinality among all possible matchings.


v) Finally, we call M perfect if any $V \setminus V(M)$ is M -saturated.

7.3 Example

Consider the graph




Then




is a maximal matching and also a maximum matching.

Further,



is a maximal matching, but not a maximum matching.

Also,



is not a maximal matching, as $M \cup \{e\}$ is still a matching.

Finally, there are no perfect matchings as $|V(G)|$ is odd.

7.4 Remark



1) Any perfect matching is a maximum matching.

2) Any maximum matching is maximal.


3) If G has a perfect matching then $|G|$ is even.

Further, then any matching M is perfect iff it is a maximum matching iff $|M| = \frac{|V|}{2}$.

7.5 Examples

Graph G	maximal	maximum	perfect
	✓	✗	✗
	✓	✓	✗

Note that there can't be a perfect matching, even though $|G|=10$ is even, as any matching uses at most one of hg, ig and jg.



We see that while it is rather easy to decide whether a matching is maximal or perfect; things are less clear for a maximum matching. Our next goal is to develop a criterion to help us decide that, called Berge's Theorem.


7.6 Definition

Let G be a graph and M a matching for G and p a path in G .

- 1) We say that p is M -alternating if its edges alternate between edges inside and outside M .
- 2) We call p M -augmenting if it is M -alternating and its start and end vertex are distinct and both not M -saturated.

How does this relate to maximum matchings?

INTERMEZZO



Ⓐ Consider G and the matching $M = \{ca, de\}$.

1) Find an M -alternating path p .


2) Create a new matching M' by removing from M all edges in $M \cap p$ and adding all edges from $p \setminus M$.

3) Repeat this process as long as possible. What properties does your final matching M have?

Ⓑ Find a graph G of size 50, where any maximum matching has size 1.

7.7 Example


Consider the graph G with matching M below.




The path $p_1(g, c, a, d, e, i, h, j)$ is M -alternating, but not M -augmenting, its start vertex g is M -saturated.

On the other hand, the path $p_2 = (b, c, a, d, e, i, h, j)$ is:

M -augmenting. In particular, it is M -alternating. What happens if we define a new edge set M' by containing all edges of M outside of p and also exactly those edges of p which were not in M ?



For $p_1 = (g, c, a, d, e, i, h, j)$ which was M -alternating, but not M -augmenting, we obtain an edge set which is not a matching.



For the M -augmenting path $P_2 = (b, c, a, d, e, i, h, j)$ on the other hand, the set M' is indeed a matching. Moreover, it contains more edges than M .

Hence, M was not a maximum matching for G .

7.8 Theorem (Berge's Theorem, 1957)

Let G be a graph and M a matching for G . Then

M is a maximum matching iff there is no M -augmenting path in G .

Proof

\Rightarrow We proceed by contraposition. Assume M is a matching and $p = (x_1, \dots, x_n)$ an M -augmenting path. We will show that M is not maximal. If the edges of p are e_1, e_2, \dots, e_{n-1} , then as x_1 is not M -saturated, we get that $e_1 \notin M$. Its p is M -alternating also e_3, e_5, \dots , i.e. all odd numbered edges are not in M , while all even numbered edges are in M . As also x_n is not M -saturated, $e_{n-1} \notin M$, whence $n-1$ is odd and n is even. Now define $M' := M \setminus \{e_2, e_4, \dots, e_{n-2}\} \cup \{e_1, e_3, \dots, e_{n-3}, e_{n-1}\}$. Note that $|M'| = |M| + 1$. We claim that M' is a matching, proving that M was not a maximum. But this is clear, as through the change of edges only x_1 and x_n are newly M' -saturated. As they were not

M -saturated, x_1 is only the endpoint of e_i in M' and x_n only of e_{n-i} in M' . Hence M' is again a matching, larger than M .

\Leftarrow : We proceed again by contraposition. Assume M is a matching of G which is not a maximum, i.e. there is another matching M' of G with $|M'| > |M|$. We aim to find an M -augmenting path in G .

To this end, we define a subgraph $H \subseteq G$ via $V(H) = V(G)$ and $E(H) = M' \Delta M$ ($= M' \setminus M \cup M \setminus M'$).

- Note that $|M' \setminus M| = |M'| - |M' \cap M| > |M| - |M' \cap M| = |M \setminus M'|$,

whence H contains strictly more edges from M' than from M .

- Further, $\Delta(H) \leq 2$. To see that, consider $x \in V(H)$ arbitrary. Then there can be at most one edge from M containing x and also at most one other edge from M' , as M and M' are matchings. Hence, $\delta(x) \leq 2$, as desired.

- Now we know that every connected component of H either is a cycle of even length (using the same number of edges from M and M'), or a path. As $|M' \setminus M| > |M \setminus M'|$, there must be at least one connected component in H which is a path of odd length, starting and ending with an edge in M' . This yields the desired M -augmenting path in G .



7.2 Hall's Marriage Theorem

Finding matchings becomes of special interest in bipartite graphs. Historically, the questions were visualised by trying to match couples for marriage. As this does not actually give a bipartite graph, we will put ourselves into the holiday spirit and will discuss bipartite graphs, where one part represents a set of children and the other part a set of presents. We will try to help Santa and develop a criterion to decide whether he can make all the children happy.


7.9 Example

Consider the bipartite graphs below with $V(G) = X \cup Y$, where X represents a set of children and Y a set of presents. How does the problem above translate to graph theory?




We draw one edge between $x \in X$ and $y \in Y$ whenever child x would be happy. To make all children happy, we need to choose edges, such that any $x \in X$ appears as exactly one end vertex (every child gets exactly one present), but all edges are disjoint (children don't have to share a present). i.e. we need to find a matching M s.t. every $x \in X$ is M -saturated.

For our graphs we observe the following:



↳ There is a matching which makes all children happy.



↳ There are two children which both only want the first present. Hence, there is no appropriate matching.

How can we know from the graph when an appropriate matching exists? This is solved in Philip Hall's marriage theorem.

7.10 Definition

Let G be a bipartite graph with parts X and Y . We say that X is matched into Y if there is a matching M for G s.t. every $x \in X$ is M -saturated.

7.11 Remark

As above, X is matched into Y iff there is an injective function $f: X \rightarrow Y$ s.t. $\{x, f(x)\} \in E(G)$ for all $x \in X$.

7.12 Theorem (Hall's Marriage Theorem, 1935)

Let G be a bipartite graph with parts X and Y . Then X is matched into Y iff for all sets $S \subseteq X$ we have $|S| \leq |N(S)|$.

Proof

" \Rightarrow " Assume X is matched into Y , say via $f: X \rightarrow Y$. Let $S \subseteq X$ be arbitrary. As $\{x, f(x)\} \in E(G)$ for all $x \in X$, we actually get that $f(S) \subseteq N(S)$. As further f is injective, by definition we get that $|S| \leq |N(S)|$, as desired.

" \Leftarrow " Assume now that for any $S \subseteq X$ we have $|S| \leq |N(S)|$.

We want to show that there is a matching M which matches X into Y . Let M be any maximum matching. We claim that M matches X into Y .

Aiming for a contradiction, assume not, i.e. there is some $u \in X$ which is not M -saturated. Define the set

$A := \{v \in V(G) \mid \text{exists an } M\text{-alternating } uv\text{-path}\}$. We claim that $S := A \cap X$ violates the assumption, i.e. $|S| > |N(S)|$.

We will split the proof into 2 parts. Let $T := A \cap Y$.

Let's start with some observations.

First note that as u is not M -saturated, for any M -alternating path $p = (u=u_1, u_2, \dots, u_k)$ we have

$$u_i, u_{i+1} \in M \text{ iff } i \in \mathbb{Z} \text{ iff } u_i \in T \text{ iff } u_{i+1} \in S.$$

Further, as M is maximal there are no M -augmenting paths.

In particular, any $v \in A \setminus \{u\}$ must be M -saturated.

Claim 1: $|S| - 1 = |T|$.

↳ We define a function $f: S \setminus \{u\} \rightarrow T$ via the following:

If $x \in S \setminus \{u\}$ then ex. $p_x = (u=u_1, u_2, \dots, u_k=x)$ M -alternating.

Then as $x \in S$, $u_{k-1}, u_k \in M$ and M is a matching, u_{k-1} is the only neighbour of x s.t. $u_{k-1}, x \in M$. Further $u_{k-1} \in T$ and we set $f(x) := u_{k-1}$. As M is a matching, f is injective.

Further, for any $y \in T$ by definition there is an M -alternating uy -path $p_y = (u=y_1, y_2, \dots, y_l=y)$. As $y_{l-1}, y_l \notin M$ and

y_c is M -saturated, there must be a vertex $x \in X$ s.t. $y_c x \in M$.

But then $py^1(x) = (u, y_2, \dots, y_c = y, x)$ is an M -alternating path

whence $x \in S$. By definition of f , we get that $f(x) = y$, whence

$y \in \text{range}(f)$. Thus, f is a bijection and $|S \setminus \{y\}| = |S| - 1 = |T|$.

Claim 2 : $N(S) = T$.

$\hookrightarrow \text{``}\subseteq\text{''}$ Let $w \in N(S)$, i.e. exists $s \in S$ s.t. $sw \in E(G)$. As $S \subseteq X$, the vertex

- w must be in y . Further, let p_s be an M -alternating ws -path.

Then by adjoining w to p_s , we obtain an M -alternating uw -path,

whence $w \in T$, as desired.

\supseteq Let $w \in T$ be arbitrary and $s := f^{-1}(t)$ with f defined as in

Claim 1. Then $s \in S \setminus \{y\}$ and $w \in N(s)$, as desired.

Conclusively, we have constructed a set $S \subseteq X$ s.t.

$$|S| = |T| + 1 = |N(S)| + 1 > |N(S)|, \text{ contradicting our assumptions.}$$

Thus, such a vertex u cannot exist, whence all vertices in X are

M -saturated and M matches X into Y . □

Application - System of Representatives

7.13 Definition

Let $F = \{S_1, S_2, \dots, S_k\}$ be a family of non-empty sets.

A **system of distinct representatives** for F is a set $\{x_1, x_2, \dots, x_k\}$


s.t. $x_i \in S_i$ and the x_i are pairwise distinct.

7.14 Example

Let $S_1 = \{2, 8\}$, $S_2 = \{8\}$, $S_3 = \{5, 7\}$, $S_4 = \{2, 4, 8\}$, $S_5 = \{2, 4\}$

Can we find a system of representatives for $F = \{S_1, \dots, S_5\}$?

Let's visualise the problem:



With this visualisation the question translates into asking whether F is matched into $\cup_{S \in F} S$.

Let $S = \{S_1, S_2, S_3, S_4\}$. Then $|S| = 4$. On the other hand, $|N(S)| = |\{2, 4, 8\}| = 3$. So, $|S| > |N(S)|$ and by Hall's marriage theorem, there is no system of distinct representatives for F . If we consider $F_0 := \{S_1, S_2, S_3, S_4\}$ however, then a system of distinct representatives is given by $\{2, 8, 5, 4\}$.

7.15 Theorem

Let $F = \{S_1, S_2, \dots, S_k\}$ be a family of nonempty sets. Then F has a system of representatives iff for any $I \subseteq \{1, 2, \dots, k\}$ we have that $|I| \leq |\bigcup_{i \in I} S_i|$.

Proof

Exercise, easy consequence from Hall's marriage theorem. \square

The König-Egerváry Theorem

We want to finish the chapter on matchings by relating them to yet another very important graph concept - the one of vertex covers.

7.16 Definition

Let G be a graph. A **vertex cover** for G is a vertex set $C \subseteq V(G)$ s.t. every edge of G has at least one endvertex in C , i.e. $\forall e \in E(G) \exists x \in C \text{ s.t. } x \in e$.

A vertex cover is called a **minimum vertex cover** if there is no vertex cover of smaller cardinality.

7.17 Application

Imagine a museum with many galleries. We want to position guards within the galleries that have all art pieces in sight. If we model galleries as edges and places where at least two galleries meet as vertices, then we obtain a graph. Now every vertex cover for G would provide an appropriate list of locations to place our guards. Of course, we want to minimize our spending when usually we are interested in finding a minimum vertex cover.


7.18 Remark

- 1) Every graph G has a vertex cover, namely V_G .
- 2) If C is a vertex cover for G and $D \subseteq C$, then D is a vertex cover for G .

7.19 Example

Consider G given by



Then one vertex cover



is given by $C_1 := \{a, b, c, d\}$. But also, $C_2 := \{a, e, d\}$ is a vertex cover of smaller cardinality. It is not hard to see that there is no vertex cover containing only two vertices, whence G is minimal.

INTERMEDIATE

For each of the graphs below, find a maximum matching M and a minimum vertex cover C . Can you conjecture a relation between $|M|$ and $|C|$ for arbitrary graphs G ?



We see that in G_1 , $|M| = 3 < 4 = |C|$, while in G_2 $|M| = 3 = |C|$.

What is the difference in both graphs? We note that G is bipartite.

7.20 Lemma

Let G be a graph. Many matching for G and C any vertex cover for G . Then $|M| \leq |C|$. In particular, a maximum matching contains at most as many edges as a minimal vertex cover contains vertices.

Proof

Let M be any matching for G , say $M = \{e_1, e_2, \dots, e_b\}$, and C any vertex cover. For any $e_i \in M$ there is a vertex $x_i \in C$ incident with e_i . As all e_i 's are disjoint, all the x_1, x_2, \dots, x_b are distinct. Thus, C contains at least as many elements as M , i.e. $|M| \leq |C|$. □

Now, what changes if we restrict ourselves to bipartite graphs?

Somehow we have more control over the relation between vertices and edges. Indeed, the Hungarian mathematicians Dénes König and Jenő Egerváry independently discovered the following in 1931.

7.21 Theorem (König - Egerváry)

Let G be a bipartite graph. Then any maximum matching has the same cardinality as any minimum vertex cover.

Proof

Consider an arbitrary bipartite graph G and a maximum matching M .

We will show that there exists a vertex cover C s.t. $|M| = |C|$.

By Lemma 7.20, C then is a minimum vertex cover. Let X and Y form a bipartition of V_G .

If every $x \in X$ is M -saturated, then $|X| = |M|$. Clearly, as G is bipartite, the set $C := X$ is a vertex cover for G , whence $|C| = |X| = |M|$ is as desired.

Now, assume not every $x \in X$ is M -saturated.

Let $U := \{x \in X \mid x \text{ is not } M\text{-saturated}\}$. Then $|M| + |U| = |X|$. (*)

Similar to Hall's Lemma, set

$A := \{v \in V(G) \mid \text{ex. an } M\text{-alternating } uv\text{-path for some } u \in U\}$.

Further, set $S := A \cap X$ and $T := A \cap Y$. We claim that

$C := (X \setminus S) \cup T$ is a vertex cover with $|C| = |M|$.

Exactly as in the proof of 7.12, we can show that

1) Every vertex in $(S \setminus U) \cup T$ is M -saturated.

2) $|S \setminus U| = |T|$ and

3) $T = N(S)$.

Thus, we immediately get that $|T| = |S| - |U|$ (**) and thus

$$\begin{aligned} |C| &= |X \setminus S| \cup |T| && \stackrel{X \cap T = \emptyset}{=} |X \setminus S| + |T| \\ &&& \stackrel{S \subseteq X}{=} |X| - |S| + |T| \\ &&& \stackrel{(**)}{=} |X| - |S| + |S| - |U| \\ &&& = |X| - |U| \stackrel{(*)}{=} |M|, \text{ as desired.} \end{aligned}$$

It remains to show that C is a vertex cover for G .

To this end, let $e = xy$ be an arbitrary edge with $x \in X, y \in Y$. We need

to show that at least one of x or y is in $C = (X \setminus S) \cup T$.

If $x \notin S$, i.e. $x \in X \setminus S$, we are done. Otherwise $x \in S$, whence

$y \in N(S) = T$, as desired. This finishes the proof. □

8. Ramsey Theory

"Complete Disorder is impossible"

8.1 The party problem

Imagine you want to plan a party and you want to avoid that a few guests only will end up alone. You want to know how many guests you will have to invite such that either m many mutually know each other or n -many are mutual strangers. We will see that this will always happen eventually. That number is called the Ramsey number $R(m,n)$.

8.2 History

- idea based on a paper by Frank Ramsey "On a problem of formal logic", (1928).
- Ramsey (1903 - 1930), although dying very young, contributed to many different areas of research. - Economy (Ramsey pricing), philosophy (Ramsey sentences), logic (Ramsey expansions) and mathematics (Ramsey numbers).



8.3 Definition

Let G be a graph. An **edge k -colouring** of G is any function $K : E_G \rightarrow \{1, 2, \dots, k\}$. In this course, we will mainly consider 2-colourings of edges and set the colours to be {red, green}.

Note that there are no conditions on the function except for the domain and codomain.

8.4 Example

Consider the following edge colourings of K_5 .



Note that K produced a triangle on $\{a, b, d\}$'s whose edges are all red, whereas with K' there is neither a red nor a green triangle.

8.5 Definition

- 1) Let G be a graph and $K: E_G \rightarrow \{1, 2, \dots, k\}$ an edge colouring. We call a subgraph $H \subseteq G$ **monochromatic** if all edges of H have the same colour, i.e. $K|_H : E_H \rightarrow \{1, \dots, k\}$ has a range of size 1, $|K(E_H)| = |\{i\}| = 1$. We then call H a graph of colour i .
- 2) The **Ramsey number $R(a, b)$** for $a, b \in \mathbb{Z}_+$ is the smallest integer n s.t. any edge 2-colouring of K_n either contains a red K_a or a green K_b as a subgraph.

8.6 Example

- 1) First note that $R(a, b) = R(b, a)$ for all $a, b \in \mathbb{Z}_+$. (Exercise)
- 2) Let's observe $R(1, b)$ is the least $n \in \mathbb{Z}_+$ s.t. any edge 2-colouring contains either a red K_1 or a green K_b . As K_1 does not contain any edges, every edge colouring of any non-empty graph satisfies the condition, whence $R(1, b) = 1$.
- 3) $R(2, 2) = 2$, as any edge 2-colouring of K_2 either colours the unique edge red or green, whence we either obtain a red or green K_2 .

INTERMÉDIO

Try to find $R(2, m)$ for an arbitrary $m \in \mathbb{Z}_+$. Prove your answer!

8.7 Lemma

Let $r \geq R(a, b)$ for $a, b \in \mathbb{Z}_+$. Then any edge 2-colouring of K_r produces either a red K_a or a green K_b .

Proof

Let $k := R(a, b)$, $r \geq k$ and \mathcal{K} be any edge 2-colouring of K_r . Clearly, K_r contains K_k as a subgraph. Then K_k , and hence K_r , either contains a red K_a or a green K_b , as desired. \square


8.8 Lemma

We have that $R(3, 4) = 9$.

Proof

We have to show two things : 1) $R(3, 4) > 8$ and 2) $R(3, 4) \leq 9$.

1) $R(3, 4) > 8$, i.e. exists an edge 2-colouring of K_8 without a red K_3 and green K_4 :




2) $R(3, 4) \leq 9$, i.e. any edge 2-colouring of K_9 either produces a red K_3 or a green K_4 .

Claim: There is a vertex v with either at least 4 red or at least 6 green edges incident.

Otherwise, as all vertices have degree 8, all vertices must be incident with exactly 3 red and 5 green edges. Consider then the subgraph $H \subseteq G$ consisting of all 9 vertices of G and all red edges. Then H is a graph which has odd many (i.e. 9) vertices of odd (i.e. 3) degree, contradicting the Handshaking Lemma.

Now consider first the case that ex. $v \in V_G$ incident with 4 red edges, say $\{vx_1, vx_2, vx_3, vx_4\}$. If any of the edges $x_i x_j$ is also red, we have a red K_3 on $\{vx_i x_j\}$. Otherwise, all these edges are blue and we get a blue K_4 on $\{x_1 x_2 x_3 x_4\}$, as desired.

Picture:



Now, in the leftover case, v has six green incident edges, say $\{vx_1, vx_2, vx_3, vx_4, vx_5, vx_6\}$. As $R(3,3)=6$ (HW), we get that there is either a red K_3 contained in $\{x_1, x_2, \dots, x_6\}$ or a green K_3 , say on $\{x_i, x_j, x_k\}$. But then we find a green K_4 on $\{v, x_i, x_j, x_k\}$ and again our claim holds.

We thus saw that there is a 2-colouring of the edges of K_8 without a red K_3 or green K_5 , however every 2-colouring of the edges of K_9 produces either a red K_3 or a green K_4 . Thus, $R(3,4)=9$.

8.9 Summary and more...

Let us sum up some of the known Ramsey numbers.

a	b	R(a,b)	a	b	R(a,b)
1	k	1	3	7	23
2	k	k	3	8	28
3	3	6	3	9	36
3	4	9	4	4	18
3	5	14	4	5	25
3	6	18	5	5	???

The problem gets very fast very complicated. Why should there even always be an answer?

8.10 Theorem (Ramsey)

For any positive integers a, b the Ramsey number $R(a, b)$ exists, i.e. there is a positive integer n s.t. every edge 2-colouring of K_n either produces a red K_a or a green K_b .

Proof

We establish the proof by induction on $m := a+b$.

$m=2$: If $a+b=2$, then $a=b=1$ and we know that $R(1,1)=1$.

$m \rightarrow m+1$: Assume for any $\tilde{a}, \tilde{b} \in \mathbb{Z}_+$ with $\tilde{a}+\tilde{b} \leq m$ we know that $R(\tilde{a}, \tilde{b})$ exists. Now assume $a, b \in \mathbb{Z}_+$ s.t. $a+b=m+1$. Then both $R(a-1, b)$ and $R(a, b-1)$ exist by I.H. Set $n := R(a-1, b) + R(a, b-1)$ and consider K_n . We claim that K_n does the job.

To this end, pick an arbitrary edge 2-colouring K of K_n and choose $v \in V(K_n)$ arbitrary. Define two sets:

$$A := \{w \in V(K_n) \mid K(vw) = \text{red}\} \text{ and } B := \{w \in V(K_n) \mid K(vw) = \text{green}\}.$$

We know that $A \cup B \cup \{v\} = V(K_n)$. and A, B are disjoint.

$$\text{Thus, } |A| + |B| = |V(K_n)| - 1 = n - 1 = R(a-1, b) + R(a, b-1) - 1.$$

$$\text{Thus either (1) } |A| \geq R(a-1, b) \text{ or (2) } |B| \geq R(a, b-1).$$

If (1) $|A| \geq R(a-1, b)$, then there exists either a green K_b or a red K_{a-1} in A , say on $X \subseteq A$. But then either X is a green K_b in K_n or $X \cup \{v\}$ is a red K_a in K_n as desired.

Similarly, if (2) $|B| \geq R(a, b-1)$, then there exists either a red K_a or a green K_{b-1} in B , say on $Y \subseteq B$. But then either Y is a red K_a in K_n or $Y \cup \{v\}$ is a green K_b in K_n , as desired.

Thus, in any case we find either a red K_a or a green K_b in K_n , whence the Ramsey number $R(a, b)$ exists and is bounded by

$$R(a, b) \leq R(a-1, b) + R(a, b-1).$$



So we saw that Ramsey numbers always exist whence it should not be too hard to find them, right? WRONG. The list given in 8.9 actually lists all currently known Ramsey numbers and even though there is continuous work in the area and bounds keep improving, the problem stays very hard.

This is nicely illustrated by the following quote by Erdős:

"Suppose aliens invade the earth and threaten to obliterate it in a year's time unless human beings can find the Ramsey number $R(5, 5)$. We could marshal the world's best minds and fastest computers, and within a year we could probably calculate the value. If the aliens demand $R(6, 6)$ however, we would have no choice but to launch a preemptive attack."

8.2 Excursion into Graph Ramsey Theory

8.11 Definition

Let G, H be arbitrary graphs. Then Ramsey number $R(G, H)$ is the smallest $n \in \mathbb{Z}_+$ s.t. every edge 2-colouring of K_n contains either a red copy of G or a green copy of H as a subgraph.

8.12 Lemma

For any graphs G, H we have $R(G, H) \leq R(|G|, |H|)$. In particular, the Ramsey number $R(G, H)$ always exists.

Proof

Let G, H be given and let $n = R(|G|, |H|)$. Consider any edge 2-colouring of K_n . Then it either contains a red copy of $K_{|G|}$ or a green copy of $K_{|H|}$. But as $G \subseteq K_{|G|}$ and $H \subseteq K_{|H|}$, we are done. \blacksquare

8.13 Example

We claim that $R(C_3, P_3) = 5$, where P_3 is the path on 3 vertices.

First, note that $R(C_3, P_3) > 4$, as  is a colouring of K_4 which neither contains a red C_3 nor a green P_3 .


Now consider an arbitrary edge 2-colouring of K_5 . Pick $v_i \in V(K_5)$ arbitrary.

If two of the edges incident with v_i are green, we obtain a green P_3 .


Otherwise, all but one edge must be red, say $v_i v_1, v_i v_2$ and $v_i v_4$ are red.

Following the same argument, if both of v_2v_3 and v_2v_4 are green, we obtain a green P_3 . Otherwise, at least one, say v_2v_3 , must be red and we obtain a red G on $\{v_1, v_2, v_3\}$.


Picture:



$$(v_4, v_1, v_5) = P_3$$



$$(v_3, v_2, v_4) = P_3$$



$$(v_1, v_2, v_3) = C_3$$