

Chapter 02

Mostafa Touny

September 13, 2023

Contents

| Problems | |
|----------|---|
| 1 | 2 |
| 2 | 2 |
| 3 | 2 |
| 5 | 2 |
| 7 | 2 |
| 14 | 2 |
| 16 | 3 |
| 18 | 3 |
| 33 | 4 |
| 34 | 4 |
| 47 | 5 |

$$\frac{13}{15} = \frac{8.7}{10}$$

Problems

1

(b). No. $\frac{3}{2}$ is not an integer. ✓ 



(d). Yes. cA is a totally valid matrix for any scalar c or matrix A .  Matrix multiplication

2

(a). Yes.  $(a - b) - c$

$$F \quad a - (b - c) = (a - b) + c$$

(b). No. $\frac{1/2}{3} = \frac{1}{2} \cdot \frac{1}{3} \neq \frac{3}{2} = \frac{1}{2/3}$. 

$$\frac{1}{1 \cdot 3}$$

(e). No. $(2^2)^3 = 2^6 \neq 2^8 = 2^{(2^3)}$. 

3

(c). No. $3(x^2) \neq 3^2x^2 = (3x)^2$. 

$$\textcircled{1}$$

(d). No. Known from linear algebra.  (An example would be better)

5

(a). $20 - 13 = 7$. 

(b). The problem is reduced to finding x and y such that $13x = 14y + 1$. In other familiar notation from chapter 1, $13x - 14y = 1$. Clearly $13(-1) + (-14)(-1) = 1$ so $13(-1 + 14) + (-14)(-1 + 13) = 1$. Thus the inverse of 13 is 13. 



7

Not closed. $1 + 3 = 4$. 

$$\textcircled{\frac{1}{2}}$$

No inverse. $3 + x \neq \boxed{1}$ for any odd integer x . 

Yes, there are inverses
for every odd number

14

$(ab)^3 = ababab$. 

$$(ab^{-2}c)^{-2} = (ab^{-2}c)^{-1}(ab^{-2}c)^{-1} = c^{-1}b^{-3}a^{-1}c^{-1}b^{-3}a^{-1} \quad \textcircled{b} \rightarrow$$

$$\textcircled{\frac{1}{2}}$$

16

Fact. x^n is an odd integer for any odd x .

Fact. The summation of two even integers is even.

~~Two even numbers added together is even.~~

We take a different perspective of the problem by the set $\{(5 \cdot 1), (5 \cdot 3), (5 \cdot 5), (5 \cdot 7)\}$ modulo $5 \cdot 8$. Upon multiplying any two elements we get the form $5 \cdot 5 \cdot x \cdot y$ where $x, y \in \{1, 3, 5, 7\}$. Think of the output of multiplication as the factor of 5 deciding the element. Observe the element is decided by $5 \cdot x \cdot y \pmod{8}$. For example if we knew $5 \cdot 5 \cdot 5 \cdot 1 = (5)(8 + 8 + 8 + 1)$ then we can easily deduce the output of $\pmod{5 \cdot 8}$ operation is $(5)(1)$.

The numbers 1, 3, 5, and 7 are all odds. So whatever x or y chosen, $5 \cdot x \cdot y$ will be odd. It follows $\text{odd} \pmod{8} = \text{odd} \in \{1, 3, 5, 7\}$. To see why note $8k + \text{odd} = \text{odd}$.

Lemma. The given set is closed under the given operation.

Lemma. The identity is $5 \cdot 5 = 25$.

Observe $5 \cdot 5 \cdot x \pmod{8} = 24x + x \pmod{8} = x \pmod{8}$ since $24x \pmod{8} = 0$.

Lemma. The inverse of $5x$ is $5x$ by computation on the given elements.

Lemma. Associativity is known from integers and modulus properties. *Show this*

2

18

$$(R_0)^2 = (R_{180})^2 = H^2 = V^2 = D^2 = (D')^2 = R_0.$$

$$(R_{90})^2 = (R_{270})^2 = R_{180}.$$

$$\text{So } K = \{R_0, R_{180}\}, \text{ and } L = \{R_0, R_{180}, H, V, D, D'\}.$$



33

Observe the group follows the same pattern as \mathbb{Z}_4 .

| | e | a | b | c | d |
|---|---|---|---|---|---|
| e | e | a | b | c | d |
| a | a | b | c | d | e |
| b | b | c | d | e | a |
| c | c | d | e | a | b |
| d | d | e | a | b | c |

inverses. Since $ad = e$, $d = a^{-1}$. Since $bc = e$, $c = b^{-1}$.

$ab = c$. $ab = (cc)b = c(cb) = ce = c$.

$db = a$. $db = d(aa) = (da)a = ea = a$.

$cd = b$. $cd = c(bb) = (cb)b = eb = b$.

$dc = b$. $dc = (bb)c = b(bc) = be = b$.

V.S
I.S

$$ac = d. \quad d = bb = (aa)(dc) = a(ad)c = ac.$$

$$bd = a. \quad bd = (dc)(bb) = d(cb)b = db = a.$$

$$dd = c. \quad dd = (ac)(bb) = a(cb)b = ab = c$$

34

(\leftarrow). Given $ab = ba$

$$\begin{aligned}(ab)^2 &= (ab)(ab) \\&= a(ba)b, \text{ Associativity} \\&= a(ab)b \\&= (aa)(bb), \text{ Associativity} \\&= a^2b^2\end{aligned}$$

(\rightarrow). Given $(ab)^2 = a^2b^2$

$$\begin{aligned}(ab)^2 &= (ab)(ab) \\&= a(ba)b \\&= aabb\end{aligned}$$

$ba = ab$, Cancellation

Once from the left
& Once from the right

(\leftarrow). Given $ab = ba$

$$\begin{aligned}(ab)^{-2} &= (ab)^{-1}(ab)^{-1} \\&= b^{-1}a^{-1}b^{-1}a^{-1} \\&= b^{-1}(ba)^{-1}a^{-1} \\&= b^{-1}(ab)^{-1}a^{-1} \\&= b^{-1}b^{-1}a^{-1}a^{-1} \\&= (b)^{-2}(a)^{-2}\end{aligned}$$

(\rightarrow). Given $(ab)^{-2} = b^{-2}a^{-2}$

$$\begin{aligned}(ab)^{-1}(ab)^{-1} &= b^{-1}a^{-1}b^{-1}a^{-1} \\&= b^{-1}b^{-1}a^{-1}a^{-1}\end{aligned}$$

$a^{-1}b^{-1} = b^{-1}a^{-1}$, Cancellation again!

$(ba)^{-1} = (ab)^{-1}$ & since inverses are unique for each element, then they must be the same!

Now observe by the definition of inverse, if $x = y^{-1}$ then $y = x^{-1}$. Therefore $ab = [(ab)^{-1}]^{-1}$ and $ba = [(ba)^{-1}]^{-1}$, and $ab = ba$.



47

Clearly $aabb = a^2b^2 = ee = e$, and $abab = (ab)^2 = e$. It follows $aabb = abab$, and by cancellation $ab = ba$.

~~2~~
①
1